











Hamilton James <sup>87.597</sup>  
from  
Uncle Nigel

March 1908.

u  
What we think: we become."







NESTS AND EGGS OF BRITISH BIRDS.

---

IN THREE VOLUMES.—VOLUME III.







5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100



PLATE  
CLVII



NUMIDIAN CRANE.  
CRANE.


CLVII



A  
NATURAL HISTORY  
OF THE  
NESTS AND EGGS  
OF  
BRITISH BIRDS

THIRD EDITION

NEWLY REVISED, CORRECTED AND ENLARGED



BY

THE REV. F. O. MORRIS, B.A.

*Rector of Nunburnholme, Yorkshire*

HONORARY MEMBER OF THE YORKSHIRE PHILOSOPHICAL SOCIETY

IN THREE VOLUMES

VOLUME THE THIRD

WITH TWO HUNDRED AND FORTY-EIGHT COLOURED PLATES

LONDON

JOHN C. NIMMO

14, KING WILLIAM STREET, STRAND

MDCCCXCII



QL675

M6

1892

v. 3

BIOLOGY  
LIBRARY  
G

TO THE  
BIOLOGY LIBRARY



# CONTENTS OF THE THIRD VOLUME.

	PAGE		PAGE
CRANE . . . . .	1	BROWN SNIPE . . . . .	39
NUMIDIAN CRANE . . . . .	2	CURLEW SANDPIPER . . . . .	40
HERON . . . . .	3	KNOT . . . . .	41
PURPLE HERON . . . . .	4	BARTRAM'S SANDPIPER . . . . .	42
GREAT WHITE HERON . . . . .	6	YELLOW-SHANKS . . . . .	43
LITTLE EGRET . . . . .	7	BUFF-BREASTED SANDPIPER . . . . .	43
BUFF-BACKED HERON . . . . .	8	BROAD-BILLED SANDPIPER . . . . .	44
SQUACCO HERON . . . . .	9	STINT . . . . .	45
NIGHT HERON . . . . .	10	PEEP . . . . .	45
BITTERN . . . . .	11	TEMMINCK'S STINT . . . . .	46
LITTLE BITTERN . . . . .	12	SCHINZ'S SANDPIPER . . . . .	46
AMERICAN BITTERN . . . . .	13	PECTORAL SANDPIPER . . . . .	47
WHITE STORK . . . . .	14	DUNLIN . . . . .	48
BLACK STORK . . . . .	15	PURPLE SANDPIPER . . . . .	49
SPOONBILL . . . . .	16	LAND-RAIL . . . . .	50
IBIS . . . . .	17	SPOTTED CRAKE . . . . .	51
FLAMINGO . . . . .	18	LITTLE CRAKE . . . . .	53
CURLEW . . . . .	19	BAILLON'S CRAKE . . . . .	54
ESQUIMAUX CURLEW . . . . .	20	WATER-RAIL . . . . .	55
WHIMBREL . . . . .	20	MOOR-HEN . . . . .	56
SPOTTED REDSHANK . . . . .	22	COOT . . . . .	58
REDSHANK . . . . .	23	GREY PHALAROPE . . . . .	61
GREEN SANDPIPER . . . . .	24	RED-NECKED PHALAROPE . . . . .	62
WOOD SANDPIPER . . . . .	25	GREY-LAG GOOSE . . . . .	63
COMMON SANDPIPER . . . . .	26	BEAN GOOSE . . . . .	65
SPOTTED SANDPIPER . . . . .	27	PINK-FOOTED GOOSE . . . . .	66
GREENSHANK . . . . .	28	WHITE-FRONTED GOOSE . . . . .	67
AVOCET . . . . .	29	BERNACLE GOOSE . . . . .	68
STILT . . . . .	30	BRENT GOOSE . . . . .	68
BLACK-TAILED GODWIT . . . . .	31	SNOW GOOSE . . . . .	69
BAR-TAILED GODWIT . . . . .	31	RED-BREASTED GOOSE . . . . .	69
RUFF . . . . .	33	EGYPTIAN GOOSE . . . . .	70
WOODCOCK . . . . .	34	CANADA GOOSE . . . . .	71
GREAT SNIPE . . . . .	35	BEWICK'S SWAN . . . . .	73
SNIPE . . . . .	36	MUTE SWAN . . . . .	74
JACK SNIPE . . . . .	38	SHELDUCK . . . . .	76



	PAGE		PAGE
RUDDY SHELDUCK . . . . .	77	GREAT AUK . . . . .	128
SHOVELER . . . . .	78	CORMORANT . . . . .	129
GADWALL . . . . .	79	GREEN CORMORANT . . . . .	130
PINTAIL . . . . .	80	GANNET . . . . .	131
WILD DUCK . . . . .	81	CASPIAN TERN . . . . .	132
GARGANEY . . . . .	83	SANDWICH TERN . . . . .	133
BLUE-WINGED TEAL . . . . .	85	ROSEATE TERN . . . . .	135
TEAL . . . . .	86	COMMON TERN . . . . .	136
WIGEON . . . . .	87	ARCTIC TERN . . . . .	137
AMERICAN WIGEON . . . . .	88	SOOTY TERN . . . . .	138
STELLER'S EIDER . . . . .	88	WHISKERED TERN . . . . .	139
EIDER DUCK . . . . .	89	GULL-BILLED TERN . . . . .	140
KING DUCK . . . . .	91	LESSER TERN . . . . .	141
VELVET SCOTER . . . . .	92	BLACK TERN . . . . .	142
SCOTER . . . . .	93	WHITE-WINGED BLACK TERN . . . . .	143
SURF SCOTER . . . . .	94	NODDY . . . . .	144
RED-CRESTED WHISTLING DUCK . . . . .	95	SABINE'S GULL . . . . .	145
POCHARD . . . . .	96	LITTLE GULL . . . . .	146
FERRUGINOUS DUCK . . . . .	97	BUONAPARTE'S GULL . . . . .	146
SCAUP . . . . .	98	MASKED GULL . . . . .	147
TUFTED DUCK . . . . .	99	BLACK-HEADED GULL . . . . .	148
LONG-TAILED DUCK . . . . .	100	MEDITERRANEAN BLACK-HEADED GULL . . . . .	150
HARLEQUIN DUCK . . . . .	101	GREAT BLACK-HEADED GULL . . . . .	150
GOLDEN-EYE . . . . .	102	LAUGHING GULL . . . . .	151
SMEW . . . . .	103	KITTIWAKE . . . . .	152
HOODED MERGANSER . . . . .	104	IVORY GULL . . . . .	154
RED-BREASTED MERGANSER . . . . .	105	ICELAND GULL . . . . .	155
GOOSANDER . . . . .	106	COMMON GULL . . . . .	156
GREAT CRESTED GREBE . . . . .	108	LESSER BLACK-BACKED GULL . . . . .	157
RED-NECKED GREBE . . . . .	109	HERRING GULL . . . . .	158
DUSKY GREBE . . . . .	110	GREAT BLACK-BACKED GULL . . . . .	159
EARED GREBE . . . . .	111	GLAUCCUS GULL . . . . .	160
DABCHICK . . . . .	112	SKUA . . . . .	161
GREAT NORTHERN DIVER . . . . .	114	POMERINE SKUA . . . . .	162
BLACK-THROATED DIVER . . . . .	115	RICHARDSON'S SKUA . . . . .	163
RED-THROATED DIVER . . . . .	116	LONG-TAILED SKUA . . . . .	164
GUILLEMOT . . . . .	117	FULMAR . . . . .	165
BRUNNICH'S GUILLEMOT . . . . .	119	MANX SHEARWATER . . . . .	166
RINGED GUILLEMOT . . . . .	120	BULWER'S PETREL . . . . .	167
BLACK GUILLEMOT . . . . .	121	WILSON'S PETREL . . . . .	168
ROTCHÉ . . . . .	123	FORK-TAILED PETREL . . . . .	169
PUFFIN . . . . .	124	STORM PETREL . . . . .	170
RAZOR-BILL . . . . .	126		



# NESTS AND EGGS

OF

## BRITISH BIRDS.

---

### CRANE.

COMMON CRANE.

PLATE CLVII.—FIGURE II.

*Grus cinenea,*

FLEMING. SELBY.

*Ardea grus,*

PENNANT. MONTAGU.

THE Crane nidificates both on low trees, stumps, or bushes, and also on the top of some old building, as well as upon a conglomerate mass of rushes or other water plants, among high grass or reeds, and in osier beds, and other such situations, in morasses, and by the sides of lakes. The nest is a large structure, made of sticks, with grass, rushes, flags, reeds, and other soft materials. The young remain for some days in the nest, and are fed by the parents with food prepared in their own crops.

The eggs are two—they are richly coloured, of a



pale olive-green ground, blotted and spotted with darker shades of green and olive-brown. Both birds incubate them.

---

## NUMIDIAN CRANE.

PLATE CLVII.—FIGURE I.

<i>Crus virgo</i> ,	PALLAS. TEMMINCK. BRISSON.
<i>Ardea virgo</i> ,	LINNÆUS. GMELIN. LATHAM.
<i>Anthropoides virgo</i> ,	VIELLOT. BONAPARTE.

THE nest is made of dry herbs and sticks.


The eggs are two in number. They are of a dull rather light yellowish-brown colour, marked over, mostly at the thicker end, with a somewhat darker shade.



Small, faint, illegible markings or text, possibly a stamp or signature, located in the upper right quadrant of the page.



524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000



HERON.  
PURPLE HERON.

CI.VIII



## HERON.

HERN. HERONSHAW. HERONSEWGH.  
COMMON HERON. CRESTED HERON.

PLATE CLVIII.—FIGURE I.

*Ardea cinerea*,  
*Ardea major*,  
*Ardea cristata*,

LATHAM. SHAW.  
LINNÆUS.  
BRISSON.

THE Heron builds, according to circumstances, either on the ground, in which situation Montagu saw several, or on trees of any sort; also, it is said, on cliffs, preferring situations in the vicinity of water. Many nests are often placed on it together—as many as eighty have been counted in one tree. Preparations for nidification are made about the month of April. The nest is placed on the very summit of the tree, or as close to it as the case will admit of, and also near the extremity of the branch, the size of the bird not admitting of a ready passage inwards. The nest, flat in shape, is rather small for the size of the tenants that have to inhabit it, but in some cases is much larger than in others, probably from an old one being built on. It is made of sticks and twigs, and has a lining of wool or hair, rushes, dry grasses, water-flags, straws, or any soft materials.



## PURPLE HERON.

The eggs are generally three in number, sometimes, it is said, four or five, and of a green colour. They vary in shape, some being pointed at both ends, and others only at the lower one.

Two broods are reared in the season, and both parents assist in the work of providing the young with food: the male also feeds the female while sitting. If alarmed for their young, they soar about aloft over the nests.

---

 PURPLE HERON.

CRESTED PURPLE HERON. PURPLE CRESTED HERON.  
AFRICAN HERON.

PLATE CLVIII.—FIGURE II.

<i>Ardea purpurea,</i>	LINNÆUS.
<i>Ardea Caspica,</i>	PENNANT. MONTAGU.
<i>Ardea variegata,</i>	LATHAM.
<i>Ardea Botaurus,</i>	GMELIN.
<i>Ardea stellaris major,</i>	RAY. WILLUGHBY.

WITH regard to the nidification and eggs of this bird, the following is the account given by Mr. Hewitson, as communicated to him by Mr. Hoy:—‘The Purple Heron does not begin to breed so early as the Common Heron, the end of May being the time



of incubation. They breed in society like the Common Heron, very frequently in low trees, in plantations of alder and willow, in the vicinity of rivers and large inland waters, the nests being only a few feet above the ground, upon which they are likewise sometimes placed in swamps overgrown with tall rushes, and in extensive tracts of reeds; they are large and flat, and are composed entirely of sticks, the finer towards the inside, or lined with species of dry sedge and rushes.' They are placed a few yards from the water; in shape they are flat and broad. Two or three nests are occasionally found near each other.

'The eggs are commonly four, rarely five, in number, and differ considerably in size and shape, as well as in colour.' They are generally dull green.



## GREAT WHITE HERON.

GREAT WHITE EGRET. GREAT EGRET.

PLATE CLIX.—FIGURE II.


<i>Ardea alba,</i>	PENNANT.
<i>Ardea egretta,</i>	FLEMING. TEMMINCK.
<i>Ardea candida,</i>	BRISSON.
<i>Ardea alba major,</i>	RAY.
<i>Ardea Egrettoides,</i>	GMELIN.

THE nest of this species, a very loose structure, built of sticks, reeds, flags, and rushes, and lined with dry grass or leaves, is placed generally on the ground, but sometimes on low trees, almost always, according to Audubon, overhanging the water, and many being placed in contiguity, as in the case of the common species. The same situation is resorted to, and the old nest repaired year after year, by the same pair. Low marshy places, covered with trees, by the side of ponds, lakes, and swamps, are those appropriated to nidification, and sometimes flat islands at a distance from the main land.

The male bird feeds the female while she is engaged in sitting; and when the young are hatched both parents unite in supplying their wants.

The eggs are three in number, and of a pale bluish green colour.





UNIV. OF  
CALIFORNIA

LITTLE EGRET.  
GREAT WHITE HERON.

CLIX







## LITTLE EGRET.

EGRET HERON.

PLATE CLIX.—FIGURE I.

*Ardea garzetta*,

PENNANT. MONTAGU.

THIS species, like the others, breeds in marshy places, either on the ground among reeds, or on the top of a willow stump, or on the branches of a low tree, about half-a-dozen feet from the ground. The nest is made of dry sticks and rushes, flags, reeds, and grass; the latter being placed inside.

The eggs are four or five or six, in number, and, one account says, of the same white colour as the bird, or another, of a pale bluish green colour.

The figure on the plate is from an egg in Mr. Smith's collection, which was procured by Mr. Hoy.



## BUFF-BACKED HERON.

LITTLE WHITE HERON. RED-BILLED HERON.  
RUFIOUS-BACKED EGRET.

PLATE CLX.—FIGURE I.

*Ardea æquinoctialis*,

*Ardea russata*,

*Ardea affinis*,


PENNANT. MONTAGU.

SELBY. JENVNS.

HORSFIELD.

THE figure on the plate is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.





UNIV. OF  
CALIFORNIA

BUFF-BACKED HERON.

SQUACCO HERON.

NIGHT HERON.

CLX







SQUACCO HERON.

RED-LEGGED HERON. CASTANEOUS HERON.  
 SENEGAL BITTERN. DWARF HERON. SWABIAN BITTERN.  
 SQUAIOTTA HERON.

PLATE CLX.—FIGURE II.

<i>Ardea Senegalensis</i> ,	GMELIN.
<i>Ardea Ralloides</i> ,	SCOPOLI.
<i>Ardea comata</i> ,	PALLAS.
<i>Ardea Squaiotta</i> ,	GMELIN.
<i>Ardea Castanea</i> ,	GMELIN.
<i>Ardea erythropus</i> , var. <i>B.</i> ,	LATHAM.
<i>Ardea pumila</i> ,	LATHAM.
<i>Ardea marsigli</i> ,	GMELIN.
<i>Cancrofagus rufus</i> ,	BRISSON.
<i>Cancrofagus luteus</i> ,	BRISSON.

THE nest is said to be built on trees.

The eggs appear to be, like those of others of the tribe, of a uniform rather light dull bluish green colour.



## NIGHT HERON.

NIGHT RAVEN. SPOTTED RAVEN. GARDENIAN HERON.  
 LESSER ASH-COLOURED HERON. JAMAICA NIGHT HERON.

PLATE CLX.—FIGURE III.

*Nycticorax Europæus,*

STEPHENS.

*Ardea Nycticorax,*

LINNÆUS.

*Ardea Gardeni,*

GMELIN.

*Ardea obscura,*

LATHAM.

THESE birds form heronries, either, according to Audubon, near plantations, or in the interior of retired and secluded swamps, as well as on some of the sea islands covered with evergreen trees. These are formed in low bushes, or in middle-sized or tall trees, as seems most convenient or secure. In some places the nests are placed within a few yards of the ground, many on the branches, others between the diverging boughs of the trees; in others at the tops of the trees, at an elevation of not much more than twenty feet; and in others again in tall cypresses, at a height of a hundred feet, or thereabouts. The nest is fabricated of sticks, and is large in size and flat in shape; it is lined with reeds, rushes, grass, and leaves.


The eggs are commonly four in number, sometimes five, and of a pale greenish blue colour.



卷之六



PLATE  
CLXI



LITTLE BITTERN.  
BITTERN.  
AMERICAN BITTERN.



## BITTERN.

COMMON BITTERN. MIRE DRUM. BITTER BUMP.  
BOG BUMPER.

PLATE CLXI.—FIGURE II.

*Botaurus stellaris*,  
*Ardea stellaris*,

SELBY.  
PENNANT. MONTAGU.

THE nest of this species is made of sticks, reeds, and other rough materials, and is generally placed on the ground, in the thickest part of the vegetation, not far from the water's edge. Dr. Thieneman says that the bird is careful to put it upon a mass of fallen reeds and prostrate rushes, so as to be beyond the effects of any temporary rising of the water.

The eggs, which are of a similar shape at either end, are from three to four or five in number, and of a uniform pale brown colour. The female sits on them, and the male brings her food while thus engaged. The young are soon able to shift for themselves.

The figure on the plate is from an egg (taken in Norfolk,) in Mr. Smith's cabinet.



## LITTLE BITTERN.

BOONK. RUFOUS BITTERN. LITTLE HERON.  
 LITTLE BITTERN HERON. LONG-NECK.

PLATE CLXI.—FIGURE I.

<i>Ardea minuta</i> ,	LINNÆUS. LEACH.
<i>Ardea danubialis</i> ,	GMELIN.
<i>Ardeola minuta</i> ,	PRINCE BUONAPARTE.
<i>Cancrophagus minutus</i> ,	KAUP.
<i>Butor minutus</i> ,	SWAINSON.
<i>Botaurus pusillus</i> ,	BREHM.

THE nest of this bird is generally to be found where flags grow, sometimes near, but at other times farther off from water. It is placed on hummocks in the marshes, or on strewed reeds or flags, a little above the usual rise of the water, and in some instances on the low boughs of an overhanging tree: a few have been found in bushes about a yard from the ground. It is made of such materials as the dry twigs of the willow, grass, reeds, rushes, and flags, and is a shapeless structure.

The eggs, four, and occasionally five, in number, or even six, according to Mr. Hewitson, are of a pale whitish green colour. Their incubation occupies sixteen or seventeen days.



AMERICAN BITTERN.

FRECKLED HERON. MOKOHO BITTERN. DUNKADOO.

PLATE CLXI.—FIGURE III.

<i>Ardea minor,</i>	WILSON.
<i>Ardea lentiginosa,</i>	MONTAGU. LEACH.
<i>Ardea Mokoho,</i>	VIELLOT.
<i>Ardea stellaris, var.,</i>	LATHAM.
<i>Bolauris freti Hudsonis,</i>	BRIS-ON.
<i>Butor Americana,</i>	SWAINSON.

THE nest is made in swamps.

The eggs are described by Hutchins as of a uniform dull olivaceous tint.



## WHITE STORK.

PLATE CLXII.—FIGURE I.

*Ciconia alba,*  
*Ardea ciconia,*FLEMING. SELBY.  
MONTAGU. BEWICK.

A heap of sticks and twigs, with any other coarse materials, forms the nest of this bird. It is placed on a house top, the summit of some tall chimney, the steeple of a Church, or an old tower, or turret; as well as on the highest parts of the loftiest trees, in the immediate vicinity of the most crowded thoroughfares. It becomes very large from being accumulated year after year.

The eggs are usually three or four in number, white, tinged with buff, and of a short oval form. The young are hatched after a month's incubation, and are attended to with sedulous attention by both parents, until fully fledged and able to provide for themselves. The old birds feed them from their own bills, with food they have previously swallowed.





UNIV. OF  
CALIFORNIA

WHITE STORK.  
BLACK STORK.

CLXII







## BLACK STORK.

PLATE CLXII.—FIGURE II.

*Ciconia nigra*,  
*Ardea nigra*,

FLEMING. SELBY.  
MONTAGU.

THIS species builds its nest, which is rather large, on the summit of the loftiest pine and other trees. The foundation of sticks is solidified by the addition of sods, the whole being finished with smaller twigs, rushes, feathers, hair, and all sorts of suitable lining.

The eggs, two or three, or four, or even five in number, are of a buff white colour, faintly tinged with blue.



## SPOONBILL.

## WHITE SPOONBILL.

## PLATE CLXIII.—FIGURE II.


*Phatalea leucorodia*,  
*Platalea nivea*,

PENNANT. MONTAGU.  
CUVIER,

THE nest of this bird is made in a tree, if one be situated conveniently for the purpose, or else on the stump of a willow, among rushes and weeds in water, or on the ground. It is put together of sticks, twigs, and rushes, and lined, if at all, with finer portions of the like; the former materials are made use of if it be placed in a tree. Several nests are built together when trees for the purpose are in sufficient number.

The eggs, laid in the month of May, are sometimes four, but usually two or three in number. They are white, spotted with pale reddish brown; some are without the red spots.





UNIV. OF  
CALIFORNIA

IBIS.  
SPOONBILL.  
FLAMINGO.

CLXIII







## IBIS.

GLOSSY IBIS. BAY IBIS. GREEN IBIS.

PLATE CLXIII.—FIGURE I.

*Ibis falcinellus*,

FLEMING. SELBY.

*Ibis ignea*,

STEPHENS.

*Tantalus falcinellus*,

PENNANT. MONTAGU.

*Numenius viridis*,

BRISSON.

THE nest of this species is placed among the 'debris' of reeds on some slight elevation, and is formed of dry grass, flags, and other such materials. Several nests are placed in proximity to each other.

The eggs are two or three in number, and of a pale bluish green colour. While the female is sitting, the male bird brings her food, and when not thus engaged, stands not far off, keeping sentry.

The figure on the plate is taken from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## FLAMINGO.

PLATE CLXIII.—FIGURE III.

*Phœnuopterus roseus,*

PALLAS.

THEY build, in like manner, in great numbers, together, and they manage somehow to sit on the eggs in the nest, which is built on the mud of the river side, and raised variously to a height of from four inches to some two feet, according as the situation is less or more liable to be flooded in rainy seasons.


The eggs, laid the latter end of May, are two in number, rather rough, and of a white colour.







Curlew



WHIMBREL.  
CURLEW.  
ESQUIMAUX CURLEW.



## CURLEW.

COMMON CURLEW. WHAAP. WHITTERICK.

PLATE CLXIV.—FIGURE II.

*Numenius arquata*,

PENNANT.

*Numenius major*,

STEPHENS.

*Scolopax arquata*,

MONTAGU.

THE nest of this bird, if any be made in some slight hollow, consists only of a little dry grass, twigs, or leaves, or is placed in the middle of a tuft of the former, among heather or rushes.

The eggs, laid in April and May, are four in number, and they differ much both in their ground colour and the spots. They are of a pale dull green, blotted all over with two shades of brown, and are very large for the size of the bird. They are placed 'quatrefoil' in the nest, the narrow ends inwards. The young run about almost as soon as hatched, but are not able to fly for a considerable time. Until then they are assiduously attended to by their parents. If approached, they hide themselves among the inequalities of the ground, and lie very close, the old birds endeavouring the while to attract the enemy away.



## ESQUIMAUX CURLEW.

PLATE CLXIV.—FIGURE III.

*Numenius borealis*,  
*Scolopax borealis*,

NUTTALL. AUDUBON.  
 WILSON.

THE eggs are three or four in number, and of a green colour, with a few large irregular-shaped spots of bright brown.

## WHIMBREL.

LITTLE WHAAP. WHIMBREL CURLEW.

PLATE CLXIV.—FIGURE I.

*Numenius phaeopus*,  
*Numenius Hudsonicus*,  
*Numenius minor*,  
*Scolopax phaeopus*,  
*Phaeopus arquatus*,

PENNANT.  
 BRISSON.  
 BRISSON.  
 MONTAGU.  
 STEPHENS.

THE nest of this species is a rude couch, placed among the heather on the open moor, or some hillock or low stump. A few dry grasses are the



materials of its composition, and it is scarcely hid from view.

The eggs are four in number, of a dark olive-brown colour, blotted with darker brown. They are wide at one end, and much narrower at the other, and are placed in the nest with the pointed ends inwards. They are considered good eating, and being sought for on this account, the numbers of the birds are diminished in consequence. The male and female sit on them by turns. If disturbed from them, they make great outcries to distract the intruder. The young leave the nest as soon almost as hatched, and quickly learn to skulk most warily on the approach of danger.

J. R. De Capel Wise, Esq., of Lincoln College, Oxford, has forwarded me a beautiful specimen of the egg of this bird.



## SPOTTED REDSHANK.

DUSKY SANDPIPER. BLACK-HEADED SNIPE.  
 SPOTTED SNIPE. COURLAND SNIPE. DUSKY SNIPE.

PLATE CLXV.—FIGURE I.

<i>Totanus fuscus,</i>	FLEMING. SELBY.
<i>Scolopax fuscus,</i>	MORRIS.
<i>Scolopax totanus,</i>	PENNANT. MONTAGU.

FOR the specimen of the egg, from which the drawing for the plate was made, I am indebted to the Rev. H. B. Tristram, Perpetual Curate of Castle Eden, Durham. It was taken in Finland, by Mr. Wolley, in 1854.





UNIV. OF  
CALIFORNIA

SPOTTED REDSHANK.  
REDSHANK.

CLXV







REDSHANK.

COMMON REDSHANK. REDSHANK SANDPIPER. RED-LEG.  
 RED-LEGGED HORSEMAN. RED-LEGGED SANDPIPER.  
 GAMBET SANDPIPER. STRIATED SANDPIPER.  
 SANDCOCK. POOL SNIPE.

PLATE CLXV.—FIGURE II.

<i>Scolopax calidris</i> ,	PENNANT. MONTAGU.
<i>Scolopax Totanus</i> ,	BRISSON. RAY.
<i>Totanus calidris</i> ,	FLEMING.
<i>Tringa striata</i> ,	LATHAM.
<i>Tringa Gambetta</i> ,	LINNÆUS. GMELIN. LATHAM.

THE nest of this bird, composed of a little coarse grass, is made by the marshy margins of lakes and other uncultivated watery places, on a heap of flags, or in some slight depression, or sheltered by a bush or tuft of herbage, as also, it is said, occasionally on heaths.

The eggs, deposited early in May, are pale reddish white tinged with green, and blotted, spotted, and speckled with dark red brown, most at the larger end; some varieties with bluish grey. They are four in number. The young are hatched in from fourteen to sixteen days, and immediately quit the nest under the tutelage of the female bird, the male taking no care of them; they soon are fledged, and able to provide for themselves.



## GREEN SANDPIPER.

WHISTLING SANDPIPER. WOOD SANDPIPER.

PLATE CLXVI.—FIGURE I.

*Tringa ochropus*,  
*Tringa Aldrovandi*,  
*Totanus ochropus*,

PENNANT. MONTAGU.  
 RAY.  
 FLEMING. SELBY.

THE nest of this species is either in sand on a bank, or among grass by the side of a stream.

The eggs, four in number, are of a greenish white ground colour, with dusky or dark brown and light reddish brown and grey spots, more or less dark.

The figure on the plate is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.





UNIV. OF  
CALIFORNIA

GREEN SANDPIPER.  
WOOD SANDPIPER.

CLXVI



UNIV. OF  
CALIFORNIA



## WOOD SANDPIPER.

LONG-LEGGED SANDPIPER.

PLATE CLXVI.—FIGURE II.

*Tringa glareola*,

PENNANT. MONTAGU.

*Tringa Grallatoris*,

MONTAGU.

*Totanus glareola*,

FLEMING. SELBY.

THE nest of this bird, which is extremely difficult to find, owing to the nature of the ground where it is put, is generally placed in a hollow, at but a little distance from water, among heath, or plants of the bog-myrtle, rushes, or grass. It is made of grass or other vegetable materials.

The eggs are three or four, pointed in shape, and of a pale greenish white, spotted and speckled, particularly at the larger end, with dark reddish brown. The hen bird incubates them, and her partner, watching by, rises up and hovers about any intruder.

The figure is from an egg brought from Norway, by Mr. Dann.



## COMMON SANDPIPER.

SUMMER SNIPE. SPOTTED SANDPIPER. SAND LARK.  
SAND LAVROCK.

PLATE CLXVII.—FIGURE I.

*Tringa hypoleucos*,  
*Totanus hypoleucos*,

PENNANT. MONTAGU.  
FLEMING. SELBY.

NIDIFICATION commences about the middle of April.

The nest is slight—a collection of a few leaves, a little moss, or dry grass, in a hollow in a bank, in a tuft of grass, or tussock of rushes, upon a bed of gravel, or even on a bare rock, the eggs being kept together by only a very slight inequality in the surface. It is generally thus sheltered or protected, on one side at least. It is usually built near the water's edge, but sometimes in an adjoining field, always above the highest water-mark. It is well hidden in a tuft of grass or rushes, or among the lower branches of willows and osiers, so as to be difficult to find. The same pair, if undisturbed, will return for several successive seasons to their accustomed building-place.

The eggs, four in number, are of a reddish white or cream-yellow tint, spotted and speckled with dark brown, and other marks of a lighter hue. Some are





UNIV. OF  
CALIFORNIA

COMMON SANDPIPER.  
SPOTTED SANDPIPER.

CLXVII



TO THE  
LIBRARY OF THE  
CONGRESS



of a clear very light blue ground colour, with minute brown spots all over; others with blots of deep brown. They are, as those of other waders, admirably adapted, both by their form and their position in the nest, to occupy the smallest possible degree of space, as rendered expedient by their large size in proportion to that of the bird. The young are hatched in about fourteen days, and leave the nest almost immediately. They quickly learn to hide themselves in the nearest covert, and in about a month are able to shift for themselves.

---

## SPOTTED SANDPIPER.

PLATE CLXVII.—FIGURE II.

<i>Tringa macularia</i> ,	PENNANT. MONTAGU.
<i>Totanus macularius</i> ,	SELBY. JENYNS.
<i>Totanus macularia</i> ,	TEMMINCK.

THE nest of the Spotted Sandpiper is placed in some well-hidden spot in a field, and is composed of short pieces of dry straw, or other such materials as may be furnished by the locality.

The eggs are of a pale reddish white colour, spotted and speckled with grey and brown of a darker and lighter shade. They are four in number. The young, as soon as hatched, run about with wonderful speed.

The figure on the plate is from an egg forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## GREENSHANK.

CINEREOUS GODWIT.

PLATE CLXVIII.

<i>Scolopax glottis</i> ,	PENNANT. MONTAGU.
<i>Scolopax canescens</i> ,	PENNANT. GMELIN. LATHAM.
<i>Totanus glottis</i> ,	FLEMING. SELBY.
<i>Limosa grisea</i> ,	BRISSON.
<i>Limosa glottis</i> ,	STEPHENS.

THE nest of this bird is a small hollow, with a few fragments of heath or grass placed within it.

The eggs, of a pear shape, as in the kindred species, are four in number, of a very pale yellowish-green colour, sprinkled all over with irregular spots of dark brown and blots of light purple grey, with fewest of either on the smaller end.





UNIV. OF  
CALIFORNIA

GREENSHANK.

CLXVIII




UNIV. OF  
CALIFORNIA







STILT.  
AVOCET.



STILT.  
AVOCET.

CLXIX



## AVOCET.

SCOOPER. COMMON AVOCET. SCOOPING AVOCET.  
CROOKED BILL. YELPER. COBBLER'S-AWL DUCK.

PLATE CLXIX.—FIGURE II.

<i>Recurvirostra avocetta</i> ,	FLEMING. SELBY.
<i>Recurvirostra avosetta</i> ,	PENNANT. MONTAGU.

THE nest of this species is said to be made in a hollow on some dry spot in a marsh, or on the sea bank, just above high-water mark, among the short grass, or other marine vegetation. It is lined with a little of those materials.

The eggs are described as being usually two, but sometimes three or four in number, brown or greenish white, spotted and speckled with black. The young are hatched in eighteen days, and leave the nest almost immediately. If chased, they hide themselves, with much success, among the scanty cover.

The figure on the plate is from an egg forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## STILT.

BLACK-WINGED STILT. STILT PLOVER. LONG-LEGS.  
 LONG-LEGGED PLOVER. LONG-SHANKS.  
 BLACK-WINGED LONG-SHANK.

PLATE CLXIX.—FIGURE I.

<i>Himantopus melanopterus,</i>	SELBY. JENYNS.
<i>Himantopus Plinii,</i>	FLEMING.
<i>Himantopus atropterus,</i>	MEYER.
<i>Himantopus rufipes,</i>	BECHSTEIN.
<i>Charadrius himantopus,</i>	PENNANT. MONTAGU.

A round hollow on the top of some tuft or mound serves as a receptacle for the nest of this bird.

The eggs, as represented by Professor Thieneman, are of a pale blue colour, blotted and streaked with greyish green, or olive green and dark brown.







CLXX



BLACK-TAILED GODWIT.  
BAR-TAILED GODWIT.

CLXX



## BLACK-TAILED GODWIT.

COMMON GODWIT. GODWYN. YARWHELP. YARWHIP.

LESSER GODWIT. JADREKA SNIPE.

RED GODWIT. HUDSONIAN GODWIT. SHRIEKER.

## PLATE CLXX.—FIGURE I.

<i>Limosa melanura</i> ,	LEISLER TEMMINCK.
<i>Limosa ægocephala</i> ,	FLEMING.
<i>Limosa rufa major</i> ,	BRISSON.
<i>Fedoa melanura</i> ,	STEPHENS.
<i>Scolopax Belgica</i> ,	GMELIN.
<i>Scolopax ægocephala</i> ,	LINNÆUS.
<i>Scolopax Hudsonica</i> ,	LATHAM?
<i>Scolopax limosa</i> ,	PENNANT.
<i>Scolopax Lapponica</i> ,	BEWICK
<i>Ægocephalus Bellonii</i> ,	RAY.

ABOUT the beginning of April these birds arrive at their nesting places, and begin to lay early in May, in the rough parts of swamps, and low meadows near water, the nest being composed of dry grass or other wild plants, and hidden among any coarse herbage.

The eggs are four in number, of a deep green or light olive-brown colour, faintly blotted with spots of a darker shade.

As soon as the young are able to flutter about, the old ones leave them to themselves.



## BAR-TAILED GODWIT.

COMMON GODWIT. RED GODWIT. MEYER'S GODWIT.  
 GODWYN. GREY GODWIT. RED-BREASTED SNIPE.

PLATE CLXX.—FIGURE II.

<i>Limosa rufa,</i>	BRISSON.
<i>Limosa grisea major,</i>	BRISSON.
<i>Fedoa rufa,</i>	STEPHENS.
<i>Fedoa Meyeri,</i>	STEPHENS.
<i>Fedoa pectoralis,</i>	STEPHENS.
<i>Scolopax leucophæa,</i>	LATHAM.
<i>Scolopax noveboracensis,</i>	MONTAGU.

THE egg of this species is described as of a pale yellowish-brown colour, speckled, blotted, and spotted with other darker shades of brown.


The figure on the plate is taken from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900



CLXXI  
RUFF.



RUFF.  
CLXXI



## RUFF.

REEVE, (FEMALE.) FIGHTING RUFF. SHORE SANDPIPER.  
 EQUESTRIAN SANDPIPER. YELLOW-LEGGED SANDPIPER.  
 GREENWICH SANDPIPER.

## PLATE CLXXI.

<i>Machetes pugnax,</i>	GOULD. SELBY.
<i>Tringa pugnax,</i>	PENNANT. MONTAGU.
<i>Tringa equestris,</i>	LATHAM.
<i>Tringa littorea,</i>	LINNÆUS. LATHAM.
<i>Totanus cinereus,</i>	BRISSON.

THE Reeve begins to lay the first or second week in May, and the young are hatched the beginning of June. The nest, made of coarse grass, is placed on some hillock among the same, or sedge, or rushes.

The eggs are subject to considerable variety; some are of a beautiful green ground colour, others olive brown, spotted with darker brown. They are four in number.



## WOODCOCK.

PLATE CLXXII.—FIGURE I.

*Scolopax rusticola*,

PENNANT. MONTAGU.

THE nest of this bird is built in woods and plantations, among dry grass, fern, or leaves, of which its structure is formed. It is loosely put together, but is placed in some warm spot, as, *e.g.*, at the root of a tree.

The eggs are generally four in number. They are of a yellowish white colour, blotted with pale chesnut brown. They require seventeen days incubation; the hen bird sits very close. The young run about as soon as hatched.

Two broods are hatched in the year.

Mr. D. Brown, of Trinity College, Glenalmond, Perth, has kindly forwarded me a specimen of the egg.





UNIV. OF  
CALIFORNIA

WOODCOCK.  
GREAT SNIPE.

CLXXII







GREAT SNIPE.

SOLITARY SNIPE. DOUBLE SNIPE.

PLATE CLXXII.—FIGURE II.

*Scolopax major*,  
*Gallinago major*,

PENNANT. MONTAGU.  
STEPHENS.

THE nest of the Great Snipe is placed on a tuft of grass or hillock in a marsh near to some standing water, and is lined with a little grass and fragments of other herbage.

The eggs are four in number, of a yellowish olive brown colour, spotted with two shades of reddish brown. They are hatched after an incubation of seventeen days, and then the young are tended for about a month by their parents.

The figure on the plate is from a specimen sent me by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## SNIPE.

COMMON SNIPE. WHOLE SNIPE. SNITE.  
HEATHER-BLEATER.

PLATE CLXXIII.—FIGURE I.

*Scolopax gallinago*,  
*Scolopax gallinaria*,  
*Gallinago media*,

PENNANT. MONTAGU.  
GMELIN.  
SHAW.

TOWARDS the end of March, or beginning of April, the male Snipe begins to 'call' for a mate.

The nest is commonly placed in the middle of a tuft of grass, rushes, or heather, in a shallow depression in the surface; by the side of water, as also among the heather where watery spots abound on the hill side, often at an elevation of from five hundred to a thousand feet above the level of the plain. A few chance stalks furnish its lining, if any be provided at all, which is not always the case.

The eggs are four in number, and of a very large size in proportion to that of the bird; their colour is pale yellowish, or greenish white, blotted at the larger end with two or three shades of brown. Some are spotted all over with small spots; some are of a clear green ground, some light blue, and others olive brown. Six eggs have several times been found in one nest, as





UNIV. OF  
CALIFORNIA

SNIPE.  
JACK SNIPE.

CLXXIII



TO THE  
LIBRARY OF THE  
CONGRESS



mentioned by the Rev. G. Low, in his 'Fauna Orcadensis,' but they doubtless must have been the joint contribution of two birds.

One is a bright green ground, blotted over with many brown and a few dark grey spots.

A second is a dull reddish brown ground, with large and small darker blots of the same colour, chiefly at the base, where they are run together.

A third is a bluish green ground, with markings of rust-colour, and brown, green, and grey spots and blots.

A fourth is a light greenish yellow, with fine large blots and wide waves of dark and light brown.

A fifth is a very pale green ground, most elegantly marked and striated with spots and streaks all over of brown, and a few black-hair streaks at the base.

A sixth is of a faint yellowish white ground, with a few blots of grey, and spotted over with small spots of yellowish brown.

A seventh is a yellowish white ground, blotted and spotted over, more towards the thicker end, with large yellowish brown markings.

Another is a fine yellowish green ground, with some dark greyish brown markings towards the base, the other part is spotted over with a lighter yellowish-brown.

A ninth is a very pale yellowish-green ground, with a very large rich dark brown mark towards the base, and a few spots of the same colour.

A tenth is a light reddish brown, blotted and spotted over with darker brown.



## JACK SNIPE.

JUDCOCK. HALF SNIPE.

PLATE CLXXIII.—FIGURE II.

<i>Scolopax Gallinula,</i>	PENNANT. MONTAGU.
<i>Gallinago minor,</i>	BRISSON.
<i>Gallinago minima,</i>	RAY. WILLUGHBY. STEPHENS.

THE Jack Snipe makes its nest on some grassy knoll, or in a tuft of rushes.

The eggs are four in number, and of a yellowish olive-colour, spotted about the larger end with two shades of brown.

One variety is a light yellowish-brown ground, with darker and lighter minute markings of the same colour.

A second is a light yellowish green ground, blotted and spotted over with darker yellowish-brown, brown, and grey.

A third is a light green ground, with spots and streaks of a yellowish brown, a good deal run together at the base.

J. R. De Capel Wise, Esq., of Lincoln College, Oxford, has kindly forwarded me a specimen for the use of this work.









BROWN SNIFE.

CLXXIV



BROWN SNIPE

RED-BREASTED SNIPE. GREY SNIPE. BROWN LONGBEAK.

PLATE CLXXIV.

<i>Scolopax grisea</i> ,	PENNANT. MONTAGU.
<i>Scolopax Noveboracensis</i> ,	SABINE.
<i>Macroramphus griseus</i> ,	EYTON. GOULD.

THE figure on the plate of the egg of this species is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## CURLEW SANDPIPER.

PIGMY CURLEW.

PLATE CLXXV.—FIGURE I.

*Tringa subarquata*,  
*Numenius pygmæus*,  
*Scolopax pygmæus*,

FLEMING SELBY.  
PENNANT.  
MONTAGU.

THE eggs of this bird are described as being of  
a yellowish colour, spotted with brown.





UNIV. OF  
CALIFORNIA

CURLEW SANDPIPER  
KNOT.

CLXXV







## KNOT.

RED-SANDPIPER. ASH-COLOURED SANDPIPER.

PLATE CLXXV.—FIGURE II.

*Tringa Canutus,*

MONTAGU.

*Tringa Islandica,*

PENNANT.

*Tringa cinerea,*

TEMMINCK.

*Calidris Canutus,*

FLEMING.

A tuft of grass serves as a depository for the egg of this species.

They are stated to be four in number, of a light yellowish-brown colour, marked at the larger end with grey and reddish spots, forming more or less of a belt, and least spotted towards the smaller end.

The figure on the plate is taken from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## BARTRAM'S SANDPIPER.

BARTRAM'S TATTLER.


PLATE CLXXV\*.—FIGURE I.

<i>Totanus Bartramii</i> ,	RICHARDSON AND SWAINSON.
<i>Totanus Bartramia</i> ,	TEMMINCK.
<i>Tringa Bartramia</i> ,	WILSON. AUDUBON.

THE nest is made, if at all, of a few grasses placed in a hollow scooped out of the earth.

The eggs are dull greyish yellow, with numerous spots of light purple and reddish brown.





UNIV. OF  
CALIFORNIA

BARTRAM'S SANDPIPER.  
YELLOW-SHANKS.  
BUFF-BREASTED SANDPIPER.

CLXXV \*







YELLOW-SHANKS.

YELLOW-SHANKED SANDPIPER. TATTLER.

PLATE CLXXV\*.—FIGURE II.

*Totanus flavipes,*  
*Scolopax flavipes,*

NUTTALL. AUDUBON.  
WILSON.

THE eggs appear to be of the usual type of this class of birds, the ground colour rather light dull yellowish, marked irregularly over, chiefly at the larger end, with darker and lighter reddish brown.

---

BUFF-BREASTED SANDPIPER.

PLATE CLXXV\*.—FIGURE III.

*Tinga rufescens,*

SELBY. JENYNS.

THE egg is light dull yellowish, with blots more or less dark, principally at and about the base.



## BROAD-BILLED SANDPIPER.

PLATE CLXXVI.—FIGURE I.

*Tringa platyrhyncha,*

GOULD TEMMINCK.

THIS bird is late in laying its eggs; they have been found not yet sat upon on the 24th. of June, and the young still unable to fly the last week in the following month. The nest is placed in a hummock of grass.

The eggs are, in general, of a deep chocolate-colour, sometimes spotted and mottled with a darker shade of the same. They are four in number.





UNIV. OF  
CALIFORNIA

BROAD-BILLED SANDPIPER.  
STINT.  
PEEP.

CLXXVI







## STINT.

LITTLE STINT. LITTLE SANDPIPER. BROWN SANDPIPER.  
MINUTE DUNLIN. LEAST SNIPE.

PLATE CLXXVI.—FIGURE II.

*Tringa pusilla*,  
*Pelidna minuta*,

PENNANT. MONTAGU.  
STEPHENS.

THE figure on the plate of the egg of this species is from a specimen in the collection of H. F. Walter, Esq., of Russell Square, London.

## PEEP.

LITTLE PEEP.

PLATE CLXXVI.—FIGURE III.

*Tringa pusilla*,  
*Tringa minutella*,

LINNÆUS.  
VIELLOT.

THIS, as might be supposed, is a miniature likeness, *i.e.*, a general one, of the others allied to it, the ground colour being dull yellowish, marked over more or less, chiefly at the thick end with reddish brown.



## TEMMINCK'S STINT.

TEMMINCK'S SANDPIPER. TEMMINCK'S DUNLIN.  
LITTLE SANDPIPER.

PLATE CLXXVII.—FIGURE I.

*Tringa Temminckii*,  
*Tringa pusilla*,  
*Tringa pucilla*,

SELBY.  
FLEMING.  
MONTAGU.

THE figure on the plate, of the egg of this species,  
is taken from a specimen forwarded by J. R. De  
Capel Wise, Esq., of Lincoln College, Oxford.

## SCHINZ'S SANDPIPER.

PLATE CLXXVII.—FIGURE II.

*Tringa Schinzii*,  
*Tringa Schinzii*,

BUONAPARTE. EYTON.  
GOULD. TEMMINCK

THE nest of this bird is placed by lakes or marshes  
on some grassy knoll.

The eggs are four in number, yellowish grey, or





UNIV. OF  
CALIFORNIA

TEMMINCK'S STINT.  
SCHINZ'S SANDPIPER.  
PECTORAL SANDPIPER.

CLXXVII







pale olive green, spotted with olive or chesnut brown markings.

The figure on the plate is from an egg sent me by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.

---

PECTORAL SANDPIPER.

PLATE CLXXVII.—FIGURE III.

*Tringa pectoralis*, JENYNS. EYTON. GOULD.

THE ground colour of the egg of this species is represented as dull rather light greenish, mottled and marbled over, principally about the thicker end, with more or less of reddish brown.



## DUNLIN.

PURRE. DUNLIN SANDPIPER. SEA SNIPE. STINT.  
 PLOVER'S-PAGE. LEAST SNIPE. SEA LARK.

PLATE CLXXXVIII.—FIGURE I.

*Tringa variabilis*,

*Tringa alpina*,

*Tringa cinclus*,

SELBY. JENYNS.


PENNANT. MONTAGU.

PENNANT. MONTAGU.

THE nest of the Dunlin is usually located under the shelter of some tuft or small bush, in any dry spot on moist moors and heaths, mosses or salt marshes, as well as by the sea. It is often concealed, intentionally or unintentionally, with great success, so as to be very difficult to find. Sometimes, however, it is fashioned upon the open grass which grows green and verdant here and there among the dark heather, 'lonely, lonesome, cool, and green.' A few bits of moss, withered heath, or grass, form its careless lining, if there be any in it, the same materials being for the most part merely rounded into form—a natural cradle.

The eggs are four in number, of a greenish white, greenish grey, or dull green colour, blotted and spotted with a darker and a lighter shade of brown, most so towards and at the larger end. Some have the ground a light blue inclining to dull white, others a clear





UNIV. OF  
CALIFORNIA

DUNLIN.  
PURPLE SANDPIPER.

CLXXVIII







light green richly spotted with light brown. They are deposited in the nest with the smaller ends inwards.

---

## PURPLE SANDPIPER.

SELNINGER SANDPIPER.

BLACK SANDPIPER.

SEA SANDPIPER.

PLATE CLXXVIII.—FIGURE II.

*Tringa maritima,*

PENNANT.

*Tringa Lincolnensis,*

PENNANT

*Tringa nigricans,*

MONTAGU.

*Tringa striata,*

FLEMING

THE nest of this species is fitted in some hollow in the ground with a little moss or other herbage for a lining. It is built on mountains among small pools of water, in the middle of a clump of grass.

The eggs are four in number. They are of a pyriform, *i.e.*, pear shape, of a yellowish grey colour, with small irregular spots of pale brown, crowded at the obtuse end and rare at the other. They appear to be laid at the end of June, or beginning of July.

The figure on the plate is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## LAND-RAIL.

CORN-CRAKE. DAKER HEN. MEADOW-CRAKE.

PLATE CLXXIX.—FIGURE I.

*Crex pratensis*,  
*Rallus crex*,  
*Gallinula crex*,  
*Ortygometra crex*,  
*Porphyrio rufescens*,

SELBY. JENYNS.  
 MONTAGU.  
 BEWICK.  
 FLEMING.  
 BRISSON.

THE nest of this species is placed among long grass or corn, in a furrow or some slight hollow, and is lined with a few of the leaves and stalks of the neighbouring herbage.

The eggs, commonly seven or eight, or ten, or even eleven in number, are of a pale reddish-brown, reddish, or yellowish-white colour, spotted and speckled with grey and reddish brown. They do not vary much, except in the size and greater or less number of the spots. Some are of a red-tinted ground colour, with blots of deep red brown and purple; others white, with a faint tinge of blue, and fancifully streaked and spotted all over. Two broods are wont to be reared in the year, the first being hatched between the beginning or middle of June, or later, towards the end. I have known however of a nest of ten eggs at Nunburnholme not hatched on the 19th. of July, (1876), and another of seven on the 22nd.: a young bird had





UNIV. OF  
CALIFORNIA

LAND-RAIL.  
SPOTTED CRAKE.

CLXXIX







been accidentally killed by some mowers on the 17th.

The young quit the nest when hatched, and in rather less than six weeks are able to fly. The female sits very close, and often suffers in consequence, from the unwitting scythe of the mower.

---

SPOTTED CRAKE.

SPOTTED RAIL. SPOTTED WATER-HEN. WATER-CRAKE.  
SPOTTED GALLINULE. WATER-RAIL.

PLATE CLXXIX.—FIGURE II.

*Crex porzana*,  
*Gallinula porzana*,  
*Rallus porzana*,  
*Ortygometra porzana*,

SELBY. JENYNS.  
BEWICK. FLEMING  
MONTAGU.  
STEPHENS.

THE nest of this Crake is put together of the larger and smaller stalks of rushes, reeds, and other water-plants, in marshy places, among beds of reeds, and often at the water's edge, and so arranged as to be capable of floating on the water, being thickly lined with soft grass, and the edges well rounded. Nevertheless, they are at times destroyed by the rising of floods.

The eggs are as many as from seven, eight, nine, or ten, to twelve in number, and even eighteen have been found together, but these doubtless the produce





of two birds. They are of a pale reddish white, or yellowish grey ground colour, spotted and speckled with dark reddish brown. The hen sits for three weeks to hatch the eggs. The young quit the nest very soon after being hatched.







PLATE



LITTLE CRAKE.  
BAILLON'S CRAKE.  
WATER-RAIL.

CLXXX



## LITTLE CRAKE.

OLIVACEOUS CRAKE. LITTLE GALLINULE. DWARF RAIL.

PLATE CLXXX.—FIGURE I.

*Crex pusilla*,  
*Gallinula minuta*,  
*Gallinula pusilla*,  
*Zapornia pusilla*,  
*Rallus pusillus*,

SELBY. JENYNS.  
 MONTAGU. BEWICK.  
 FLEMING.  
 GOULD.  
 LATHAM.

NIDIFICATION commences at the end of May or beginning of June.

The nest, of considerable size, is placed either in a tuft of rushes or upon the water itself, supported in the latter case on the stalks of rushes or other aquatic plants bent down and intermingled with grass, the adjoining materials of the like kind being drawn in a hood-like manner over it.

The eggs are from seven or eight to ten in number, smooth, without much polish, and of a light olive brown, yellowish, or greenish white colour, spotted with darker brown.

The young run out of the nest almost directly after being hatched.

The figure on the plate is from an egg brought from Hungary, and now in the possession of the Rev. H. B. Tristram, Perpetual Curate of Castle Eden.



## BAILLON'S CRAKE.

PLATE CLXXX.—FIGURE II.

*Crex Baillonii*,  
*Gallinula Baillonii*,  
*Zapornia Baillonii*,  
*Ortygometra Baillonii*,

SELBY. JENYNS.  
 TEMMINCK.  
 STEPHENS.  
 SHAW.

THE nest of this bird is made near the water's edge, in the moist situations which the bird itself frequents, among long grass, flags, or rushes, and is very difficult to find. Its component parts are the stems and leaves of water-plants, sedge, and grasses.

The eggs are seven or eight to ten in number, and of a regular oval form. Their colour is greyish white, spotted with yellowish brown. The hen bird is said, on leaving them, to add to the nest the concealment of the surrounding and overhanging herbage.

The figure on the plate is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## WATER-RAIL.

BILCOCK. BROOK-RUNNER. RUNNER.

PLATE CLXXX.—FIGURE III.

*Rallus aquaticus*,

PENNANT. MONTAGU.

THE nest of this species, which is built of sedge, flags, and grass, is abstrusely concealed among the thickest herbage; frequently in willow beds. Mr. Wolley mentions one which was built upon rushes, floating on the water, and another upon a clump close to its margin.

The eggs are covered with small specks of greyish ash-colour and reddish brown, on a dull cream-coloured white ground; some are almost spotless. They are from seven to ten in number.



## MOOR-HEN.

WATER-HEN. COMMON GALLINULE. MOAT-HEN.  
MOOR-COOT. MARSH-HEN.

PLATE CLXXXI.—FIGURE I.

*Gallinula chloropus*,  
*Fulica chloropus*,

PENNANT. MONTAGU.  
BEWICK. FLEMING.

THE nest of the Moor-Hen, which is large, is strongly put together, though only of rough workmanship, and is commonly found well concealed among reeds, long grass, or the roots of trees, just above the water's edge, on the margin of a stream or by a bank. It has been known as much as three feet above the surface, on the stump of a tree, or even on the lower branches of a fir, or in a thorn bush at that elevation. The Rev. Leonard Jenyns has recorded one instance in which it was constructed among the ivy encircling a large elm, which hung over the water's edge, at the height of at least ten feet from the ground. He says, 'there was a reason for it, the rising of the water in the pond frequently flooded the banks of the island, and, as I had before witnessed, had destroyed several broods.' One was built upon the branches of a willow overhanging the lake at Castle Howard, at a height of four or five feet above the water. A writer in the 'Magazine of Natural History,' mentions





UNIV. OF  
CALIFORNIA

MOOR-HEN.  
COOT.

CLXXXI







another placed in a fir tree twenty feet above the water. R. T. Davidson, Esq., of Muirhouse, has informed me of a nest he found, thirty feet from the water, near the river Blackadder, at precisely the same height in the same kind of tree.

The nest has been seen quite unattached to any fixture, though surrounded by loose sticks, and thus at the mercy of the winds and waves. Moor-Hens have been known both to hatch their eggs after being removed in part of the nest to another place, and also themselves to remove them, when threatened with destruction by the rising of the water; fresh materials being in each case brought together. J. H. Gurney, Esq. and W. R. Fisher, Esq. have recorded an instance in which they knew the nest of a Moor-Hen placed in a fir plantation at a distance of a quarter of a mile from any water. I have seen one myself placed at some height above the water on a fallen branch of a tree, which formed a natural bridge over a river.

The eggs are usually five, six, seven, or eight, in number; nine or ten have, however, been often seen in one nest. They are of a reddish or yellowish white colour, spotted and speckled all over with reddish brown; they vary exceedingly in size. Three broods are commonly reared in the year, sometimes it has been thought even four; the first eggs are laid the end of April or in May, and are, in early seasons or localities, hatched in the latter month, but otherwise not till June. It is a curious fact that the youngest brood is carefully and kindly attended to by that which is its elder, as both are at the same time by the parents, but when a third comes, it is to the abandonment of the first.

Incubation continues three weeks. The young soon leave the nest, still attended by their mother,



who leads them to the water, but, for a time, they return to it at night for shelter. The hen takes the young at times under her wings.

---

COOT.

COMMON COOT. BALD COOT.

PLATE CLXXXI.—FIGURE II.

*Fulica atra*,  
*Fulica aterrima*,

PENNANT. MONTAGU.  
LINNÆUS.

THE nest of this bird, not unfrequently deferred to be made until May, is a large structure, and, though of rough workmanship, very strong in its composition, so as to keep the eggs dry, albeit in such close proximity to water. It is built by the edges of islands in, or the borders of, lakes, ponds, and rivers, and is generally placed among, and loosely attached to, flags and reeds; sometimes on a tuft of rushes, and composed of the former plants: the finer portions are placed inwards. Mr. Hewitson says that they are sometimes accumulated so much as to rise from half a foot to a foot above the water, going down also to a depth of from one foot and a half to two feet; the width is about a foot and a half, and the interior nearly flat. Since writing the above, I have observed



one placed on the water, as indeed they not unfrequently are, and confined only in its place by the reeds springing up around it. It was three or four yards from the edge of a small pond, adjoining the high-road between Fangfoss and Stamford Bridge. The old bird moved a little way from it as I stopped, but did not appear shy, as she doubtless would at another time. One of them appears to remain on the site of the nest, while the other brings her articles with which to compose it.

Bishop Stanley writes thus on the subject of this part of the natural history of these birds:—‘They, too, build a simple rushy nest, but with this difference, that instead of seeking to raise it above the water, they seem to prefer it floating upon the very surface, where, of course, it is exposed to the double danger of being carried hither and thither according as the wind blows, or if interwoven with reeds or rushes close to the water, of being covered, should the waters be raised by floods. But the Coot is apparently well aware of these possibilities, and accordingly guards against them, preventing the nests being carried away, by ingeniously fastening the materials of which they are made, to the rushes or osiers near them; but at the same time, these fastenings are of such a nature, as to allow of the nests rising with the water, so that no ordinary flood would expose them to the danger of immersion. The Coot, like the Water-Hen, covers her nest, and, if not so effectually, yet with a most extraordinary rapidity. We have repeatedly watched a Coot quietly sitting on her nest; if the boat approaches, she rises, and immediately begins pecking away right and left, which she continues to do until the enemy is so near that she is compelled to decamp



for her own preservation. In this short time, however, she almost always contrives to cover her eggs; and though the nest itself remains a very conspicuous object, a careless observer might pass it as deserted and empty.' Quarrels arise among different birds sometimes at the spring of the year, for the choice of a situation.

The eggs are from six or seven, to ten or even fourteen in number, of a light dull yellowish or greenish pale brown, or stone-colour, spotted with small rust-coloured spots. If the first hatch be taken or destroyed, a second is produced, but in less numbers.


The young almost immediately leave the nest to run about, and after a few days entirely forsake it, unless the weather is unseasonable, in which case they return to it at night for a week or two, the old birds carefully tending them as long as necessary. The hen covers them with her wings.







NO. 100



GREY PHALAROPE.  
RED-NECKED PHALAROPE.



## GREY PHALAROPE.

PLATE CLXXXII.—FIGURE I.

<i>Phalaropus lobatus,</i>	PENNANT. MONTAGU.
<i>Phalaropus hyperboreus,</i>	BEWICK.
<i>Phalaropus platyrinchus,</i>	GOULD.
<i>Tringa lobata,</i>	MONTAGU.

THE nest of this species is described as slight, lined with a few grasses, or a mere formation of the natural herbage around, and placed in a hollow on some elevated spot in marshy places, but always in the vicinity of water, either the sea, or some inland lake or river.

The eggs are usually four in number, of a stone-colour, with a tinge of olive, and spots and specks of dark brown. The old birds shew great fondness for their young, and remain with them till they are well able to fly.



## RED-NECKED PHALAROPE.

RED-NECKED COOT-FOOT. RED-NECKED LOBE-FOOT.

PLATE CLXXXII.—FIGURE II.

*Phalaropus hyperboreus*,

PENNANT.

*Phalaropus fulicaria*,

MONTAGU.

*Lobipes hyperboreus*,

FLEMING.

*Lobipes hyperborea*,

SELBY

THESE birds are late in laying, only one or two eggs having been found laid by nearly the middle of June. It appears that the male bird assists the female in the task of incubation.

The nest is placed in a hollow on a small hillock, or in a tuft of grass by the edge of the water, and is deep in shape—the material dry grass. Both parents are much attached to each other, as well as to their offspring. Several nests are frequently placed in the same neighbourhood.


The eggs appear to be four in number, and of a dull olive-green colour, a good deal blotted with black brown or dusky. The young leave the nest soon after being hatched.



1854



Can. or  
Canada



GREY-LAG GOOSE.  
BEAN GOOSE.

CLXXXIII



## GREY-LAG GOOSE.

WILD-GOOSE. GREY-LAG WILD GOOSE.

PLATE CLXXXIII.—FIGURE I.

*Anser palustris,**Anser ferus,**Anas anser,**Anas ferus,*

FLEMING. SELBY.

JENYNS.

PENNANT.

MONTAGU.

THE nest of this Goose is made of grass, rushes, leaves, or dry stalks of plants, under cover of some rushes or osiers, and is well lined with feathers. It is large in size, and is located in a marsh, or by the border of a lake or inland sea. The male keeps guard near it while his partner sits.

The eggs are said to be commonly four or five in number, but to amount also to as many as twelve or fourteen, the former the produce of younger, and the latter of older birds; eight or nine the intermediate quantity. They are of a dull yellowish white, smooth and shining in appearance.

Meyer says, 'The female is very careful in covering the eggs with some of the surrounding materials, whenever she leaves the nest for a short time; and it may serve as a safe guide to persons who go in search of the eggs, that if they are uncovered they are forsaken, and are, consequently, not worth leaving in the nest.'



As soon as the Goose has laid her full number of eggs, she plucks the down off her breast, and disposes it in such a manner among the eggs, that they retain an equal temperature even at the changes of the weather, or during the short periods when she leaves the nest, once or twice a day.'

'In four weeks the young come forth, and after remaining under the mother the entire first day, are subsequently led to the water, and made to swim to some small islet, where they can hide and feed on the young blades of corn, grass, and duckweed. The gander redoubles his watchfulness on the increase of his family, and hardly ever leaves the party. On the approach of danger, the parents resort immediately to the shelter of rushes, standing corn, or long grass, attended by the whole brood; but when surprised on open ground, too far from shelter, the young lay themselves flat on the ground in some rut or hollow, and have even been known to be taken up in the hand, and carried away; but if they are near enough to the water, instinct teaches them to resort to that element for protection, where, by diving or swimming to the shelter of some cover, they may elude observation: on such occasions the parents fly round the intruders, uttering their inharmonious cries.' Yarrell says that when the hen birds begin to sit, the males leave them, and collect in flocks near or on the sea. The male and female are considered to unite for life. They return yearly to the same breeding places, arriving at them in March.

A wild Grey-lag gander is recorded to have paired with a tame goose in a farm-yard.



## BEAN GOOSE.

WILD GOOSE.

PLATE CLXXXIII.—FIGURE II.

*Anser ferus,*  
*Anas segetum,*FLEMING. SELBY.  
PENNANT. MONTAGU.

THE nest of the Bean Goose is built in heath.

The eggs are of a dull white colour, and from five to seven in number.

The young, till able to fly, hide themselves, if alarmed, in the neighbouring heather or other cover.

In confinement this species has been known to pair with the Tame Goose, the progeny partaking in some respects of the character of both parents. This is a statement of Meyer's.



## PINK-FOOTED GOOSE.

PLATE CLXXXIV.—FIGURE I.

*Anser phænicopus,*  
*Anser brachyrhynchus,*

BARTLETT.  
BAILLON.

THESE birds unite about the middle of May. Mr. J. Macgillivray has remarked that he saw them 'in pairs about the middle of the month, and they had the young fully fledged and strong upon the wing about the end of July.' They had again collected into flocks by the beginning of August.

The eggs are of a pure white colour. Eight were laid by one of these Geese, kept on the water in St. James's Park by the Ornithological Society of London.





UNIV. OF  
CALIFORNIA

PINK-FOOTED GOOSE.  
WHITE-FRONTED GOOSE.

CLXXXIV







WHITE-FRONTED GOOSE.

LAUGHING GOOSE. WHITE-FRONTED WILD GOOSE.

PLATE CLXXXIV.—FIGURE II.

*Anser erythropus*,  
*Anser albifrons*,

FLEMING. SELBY.  
PENNANT. MONTAGU.

THE eggs of this species are of a white colour, with  
a tinge of pale buff.



## BERNACLE GOOSE.

BARNACLE GOOSE. COMMON BERNICLE GOOSE.

PLATE CLXXXV.—FIGURE I.

*Anser bernicla*,

FLEMING. SELBY.

*Anser leucopsis*,

JENYNS.

*Anas erythropus*,

PENNANT. MONTAGU.

*Anas leucopsis*,

TEMMINCK.

THESE birds appear to breed in large swamps and morasses.

The eggs are of a greenish white colour.

## BRENT GOOSE.

PLATE CLXXXV.—FIGURE II.

*Anser brenta*,

FLEMING. SELBY.

*Anser torquatus*,

JENYNS.

*Anser bernicla*,

PENNANT. MONTAGU.

THE nest of the Brent Goose is formed of vegetable materials, collected together in swampy places.

The eggs are white, with a faint tinge of brown or greyish.





UNIV. OF  
CALIFORNIA

BERNACLE GOOSE.

BRENT GOOSE.

CLXXXV



TO THE  
LIBRARY OF THE  
MUSEUM OF NATURAL HISTORY







egg of  
Canada



SNOW GOOSE.  
RED-BREASTED GOOSE.

CLXXXV\*



SNOW GOOSE.

PLATE CLXXXV\*.—FIGURE I.

*Chen Hyperboreus,*

PALLAS.

THE nest is placed in the vicinity of lakes in a slight hollow made in the sandy soil.

The eggs, commonly five in number, are rough, and of a dull white colour.

---

RED-BREASTED GOOSE.

SIBERIAN GOOSE. RED-BREASTED BERNICLE.

PLATE CLXXXV\*.—FIGURE II.

*Anser ruficollis,*

FLEMING. SELBY.

THE eggs are like those of the other kinds, dull white all over.



## EGYPTIAN GOOSE.

PLATE CLXXXVI.—FIGURE I.

*Anser Egyptiacus,*

JENYNS.

*Anas Egyptiaca,*

BEWICK.

*Chenalopex Egyptiaca,*


GOULD.

THESE birds are believed to pair for life, and are much attached to each other, as also to their young. They make their nest, it is said, of reeds, leaves, and the stems of water-plants, and give it a lining of cotton and feathers. It is always placed near the water, and has been known on its surface, supported by the surrounding floating vegetation.

The eggs are of a dull white colour, with a tinge of buff, six or eight in number.

The figure on the plate is from one laid in the Gardens of the Zoological Society.





UNIV. OF  
CALIFORNIA

EGYPTIAN GOOSE.  
CANADA GOOSE.

CLXXXVI



TO THE  
LIBRARY OF  
CALIFORNIA



## CANADA GOOSE.

CRAVAT GOOSE.

PLATE CLXXXVI.—FIGURE II.

*Anser Canadensis*,  
*Anas Canadensis*,  
*Cygnus Canadensis*,

FLEMING.  
BEWICK.  
JENYNS.

THE Canada Goose makes for herself, or rather for her expected brood, a 'downy bed' of feathers, within a structure of dry plants of different kinds. It is of large size, and raised to the height of several inches. It is begun to be built in March or April, and is placed among deep grass, near the water. Audubon mentions his having found one on the stump of a large tree, standing in the middle of a small pond, and at a height of about twenty feet. Occasionally they build in the old nests of other large birds; and while the hen sits her consort keeps sailing about over the water in the neighbourhood, but not coming very close to the nest. On the appearance of any danger he exhibits much anxiety for his family.

The eggs are sat upon by the end of May or the beginning of June. They are generally six or seven in number, but as many as nine have been found: they are of a dull white colour.

The figure on the plate is from a specimen laid in St. James's Park.



## HOOPER.

WHISTLING SWAN. HOOPING SWAN. ELK.

PLATE CLXXXVII.

*Gygis ferus,*  
*Anas cygnus,*RAY. WILLUGHBY.  
LINNÆUS. GMELIN.

THE nest, which is of large size, as might naturally be looked for, is made of reeds, rushes, and other water-plants, and is lined with down, with which the eggs are also covered. It is about a foot and a half in diameter, and is placed not far from the water. It is begun to be made about the middle of March, and is built on the ground in marshy places.

The eggs are of a pale dull brownish white or pale greenish white colour. They vary in number from five to seven.

The female sits, it is stated, for forty-two days. She and her mate are united, it is said, for life. They keep the place of their nest free from intrusion, and resolutely repel its appropriation by any but themselves.





HOOPER.

CLXXXVII

UNIV. OF  
CALIFORNIA







20 MAY 1964



UNIV. OF  
CALIFORNIA



BEWICK'S SWAN.

MUTE SWAN.

CLXXXVIII



## BEWICK'S SWAN.

PLATE CLXXXVIII.—FIGURE I.

*Cygnus Bewickii,*

YARRELL.

THE nest is reported by Captain Lyons, R.N., to be built of peat-moss, and to be nearly as much as six feet long, four feet and three quarters wide, and two feet high on the outside, the hollow one foot and a half across. Another account says that the materials used are flags, rushes, and the small boughs of willow trees; doubtless those most readily procurable are differently made use of in different places.

The eggs are of a white cream colour. This Swan too is said to occupy six weeks in the incubation of them. Many pairs build in the same vicinity, but each pair maintain the right of private property for the time being in their own more immediate abode.



## MUTE SWAN.

TAME SWAN.

PLATE CLXXXVIII.—FIGURE II.

*Cygnus olor*,

JENYNS.

*Cygnus mansuetus*,

GOULD.

*Anas olor*,

PENNANT. BEWICK.

THE Swan disposes its nest on the ground, near the water side, or on some mound on an island in the river or lake. It is made of rushes and flags, and if the water threatens to rise, more materials, which the male bird brings, and the female works in, are added to the deposit under the eggs, which are thus gradually raised further out of danger.

The ordinary number of eggs laid by this Swan, is from two to four, sometimes five, occasionally six, and not very rarely eight in number, older birds laying the larger, and younger the fewer numbers respectively. They are of a dull greenish white colour. In one instance as many as nine were laid, and all of them successfully hatched, in the Jephson Gardens, at Leamington. It is possible, however, that two birds may have made use of the same nest, and, if so, the one under whose care all came, proved no 'injusta noverca,' but tended all with the like attention. At Beddington Park, in Norfolk, twelve eggs were deposited, and the brood all reared, in 1850.



Incubation continues for from five to six weeks. After being hatched for one day, the cygnets follow the guidance of their parents to the water, and have but little instruction beyond that instinctively given them by nature, in the art of swimming about and feeding themselves. Still, 'The attention,' says Meyer, 'bestowed by the old birds upon the young is incessant; and when fatigued by the strength of the stream, or requiring to be removed to a far distance, too great for their young capacity, the hen bird takes the young ones on her back, which she accomplishes by lowering herself a little in the water, and occasionally assisting them to ascend with her foot, and in this manner they are carried in safety to some more desirable spot. The shape of the Swan's back, which is very flat, is well adapted for this purpose; and when her wings are raised, the young ones repose in the most beautiful and safe cradle imaginable.'

The nest, when the first egg is laid, is small in size, but, as by degrees a larger family is expected, the mistress adds to the size of it by clutching at every suitable material in its vicinity, and this even to a greater extent than appears to be, or indeed is, at all necessary. Instinct suggests this for a wise purpose; but where reason would say 'hold, enough,' the former displays its inferiority by not knowing where to stop.



## SHELDUCK.

SHIELDRAKE. COMMON SHIELDRAKE. BURROW DUCK.  
 SKELGOOSE. SKEELING GOOSE.

PLATE CLXXXIX.—FIGURE I.

*Tadorna Vulpanser,*  
*Tadorna Bellonii,*  
*Anas Tadorna,*

FLEMING.  
 STEPHENS.  
 LINNÆUS.


THE Shieldrake builds in rabbit-burrows and other hollows in the earth, often as much as ten or twelve feet from the entrance, also in holes in rocks. Some down, plucked from their own breasts, is the lining with which the nest is fitted, the remainder being dry grass.

The eggs are ten or twelve, or even more, it is said thirteen or fourteen or even sixteen, in number; but these, in such cases, may possibly have been the produce of two birds. They are nearly perfectly white, having only a very faint tinge of green, and are smooth and shining. They are equally rounded at both ends.

The hen bird sits, as is believed, from about twenty-six to thirty days, her mate keeping watch hard by, and taking her place in the morning and evening while she picks up some food.

The young, when hatched, are either carried by their parents in their bills to the water, or soon make





UNIV. OF  
CALIFORNIA

SHIELDRAKE.  
RUDDY SHIELDRAKE.

CLXXXIX







their way thither themselves. They hide away at the approach of danger; the old birds, conscious no doubt that they are able thus best to find security, then flying off.

---

RUDDY SHELDUCK.

PLATE CLXXXIX.—FIGURE II.

*Anas rutila*,  
*Tadorna rutila*,

BEWICK.  
SELBY.

THIS species builds in holes in river banks, those which have been burrowed and deserted by animals, or such as are of natural formation.

The nest is lined with feathers of the bird itself, the remainder being any sort of stalks and sticks.

The eggs are white, and eight or nine or ten in number. They are highly polished.

The young are often carried to the water in the bill of the mother.

For the loan of the egg figured on the plate, I have to thank the Rev. H. B. Tristram, Perpetual Curate of Castle Eden. It was received by him from Dr. Thieneman, who had procured it, with others, from the Volga and the Caspian Sea.



## SHOVELER.

BROAD BILL. BLUE-WINGED SHOVELER.

COMMON SHOVELER.

RED-BREASTED SHOVELER. SHOVELER DUCK.

PLATE CXC.—FIGURE I.

*Anas clypeata*,

PENNANT. MONTAGU.

*Spathulea clypeata*,

FLEMING. SELBY.

*Rhyncaspis clypeata*,

GOULD.


THE nest of this species, built beside rivers, lakes, and other waters, or in watery places, appears to be made of grass, commingled with down from the bird itself. In some cases the bare earth or sand is scarcely covered with any materials; in others, a tuft of grass is laid in. After the female has begun to sit, she covers the eggs with down plucked from her own body.

The eggs are as many as eight, nine, ten, or twelve in number. They are of a buff white colour, with a tinge of green.

Incubation lasts three weeks. The young leave the nest almost immediately after being hatched, and repair with their mother to the water.

These birds bred in the year 1854, in the Gardens of the Zoological Society, London.





SHOVELER.  
GADWALL.  
PINTAIL.

CXC

UNIV. OF  
CALIFORNIA



TO THE  
LIBRARY OF THE  
CONGRESS



## GADWALL.

GADWALL DUCK. COMMON GADWALL.

PLATE CXC.—FIGURE II.

<i>Anas strepera</i> ,	PENNANT. MONTAGU. BEWICK.
<i>Chauliodus strepera</i> ,	SELBY. JENYNS.

THE nest of the Gadwall is placed among reeds, sedge, rushes, or other aquatic herbage, such as vegetates by the sides of pools, meres, and lakes, and in marshy places 'with verdure clad.' It is composed of dry grass, and the eggs are covered with down.

The eggs are from five to seven or eight, or, Selby says, ten or twelve in number. They are of a uniform buff white colour.

The engraving is from an egg laid in the Gardens of the Zoological Society. The birds bred there in the year 1839, and again in 1841, and in 1853 or 1854.



## PINTAIL.

COMMON PINTAIL. PINTAIL DUCK.

PLATE CXC.—FIGURE III.

<i>Anas acuta,</i>	PENNANT. MONTAGU. BEWICK.
<i>Querquedula acuta,</i>	SELBY.
<i>Dafila caudacuta,</i>	GOULD.


OF this species also, the nest is placed by the margin of, or at no great distance from, water, lakes, ponds, and seas, and is composed of grass and reeds, with a little lining of down. Some have been found in ditches, and even in standing corn: it is always well concealed.

These Ducks pair in April.

From six to eight or nine eggs are laid. The young are hatched in about twenty-three days. They at once repair to the water.

These birds have repeatedly bred in confinement. Montagu has recorded several instances of this. Lord Stanley had a female Pintail which paired with a Wigeon, and had eggs two successive seasons. The Hon. Twisleton Fiennes another, a male, which paired with a common Duck.







University of  
California



GARGANEY.  
WILD DUCK.

CXCI



## WILD DUCK.

COMMON WILD DUCK. MALLARD, (THE MALE BIRD.)

PLATE CXCI.—FIGURE II.

*Anas boschas,*

PENNANT. MONTAGU.

THE nest of this species, constructed the latter end of April, is placed, unless in a few rare exceptional instances, on the ground in a dry place, often near, but on the other hand not unfrequently at a distance from water; in some cases under a hedge, and in others in an open field, or in a wood, but under shelter of some kind. It is small in size, little more than six inches in the inner width, and regularly formed of dry grass or other vegetable materials; the lining being down, to the thickness of between two and three inches.

The greater number of these birds go north to breed; but not a few remain here and there throughout the country all the year, and build.

The eggs are usually eleven in number; sometimes ten, and even twelve; and as many as fifteen are said to have been found in one nest. They are smooth and of a very pale green colour.

The old birds are supposed to remain in pairs throughout the year, but the male leaves the female



as soon as she begins to sit. The hen covers the eggs with down and other substances if she leaves the nest for a time.

As Mr. Hewitson observes, 'We should scarcely expect to find the nest of the Wild Duck in a tree, and yet several instances have occurred in which it has chosen for itself a site thus elevated, and apparently uncongenial to its usual habits. Mr. Kute has met with a nest of this species in the grounds of Castle Howard, in a large oak tree, twenty-five feet above the ground, and fifty yards from the edge of the water. Mr. Marmaduke Tunstall speaks of one at Etchingham, in Sussex, which was built in an oak tree twenty-five feet above the ground, and contained nine eggs; and Mr. Selby says that a Wild Duck laid its eggs in the nest of a Crow, at least thirty feet from the ground.' This was at Madeley, in Staffordshire. The drake was also seen to perch on a bough near her, and occasionally in her absence sat on the nest. Others have been found at a height of ten and eighteen feet.

In Daniel's 'Rural Sports,' mention is made of the deserted nest of a Hawk in a large oak having been appropriated and repaired by a Wild Duck, and two eggs laid in it; and Montagu speaks of one built between the trunk and the boughs of a large elm tree, and of another in a willow tree overhanging some water. Meyer mentions one found by him on the stump of an old willow tree; and G. B. Clarke, Esq., in 'The Naturalist,' volume i., page 116, one built on the fragment of a broken branch of an oak about twelve feet from the ground, and a foot and a half from the trunk. Another, of which Mr. A. Foster-Melliar has informed me, containing nine eggs, was placed in a pollard willow tree, twelve feet from the



surface, and a hundred yards from any water. Another was found at Thornton Abbey, in Lincolnshire, near the top of a large ivy-covered ash tree; another in an old ruin. Sir William Jardine mentions one built on the top of a precipitous rock, and Audubon one in the middle of some corn, a mile from any water. Bishop Stanley records an instance of a nest placed on the roof of a 'lowly thatched cottage' by the water side, in the Regent's Park, London. A pair built near Nunburnholme, in May, 1876, a full half-mile from our stream or any other water. I saw one of the birds flying that way in the afternoon, and was told of the fact in the evening of the same day.

## GARGANEY.

GARGANEY DUCK. GARGANEY TEAL.

PLATE CXCI.—FIGURE I.

*Anas querquedula*,  
*Querquedula circia*,

PENNANT. MONTAGU.  
 SELBY.

THE Garganey begins to lay, as Mr. Hoy has recorded, in a communication to Mr. Hewitson, about the middle of April, or towards the end of the month, or the beginning of May.

The nest, according to the same authority, is made



of dry grass and rushes, mixed with the down of the bird itself. It is placed in low and marshy situations, among reeds and rushes, and by the borders of inland waters and rivers, among the adjoining rank herbage.

The eggs are from eight to ten or even fourteen in number—buff in colour.

Incubation continues for about twenty-one days.

The young birds, as soon as hatched, follow their mother to the nearest water.


The figure on the plate is from an egg laid in a basket by one of these birds captured in the county of Norfolk, and sent to Mr. Girdlestone.







U.S. BIRD  
CLUB



BLUE-WINGED TEAL.

CXCI \*



BLUE-WINGED TEAL.

PLATE CXCI\*.

*Querquedula Discors,*

LINNÆUS.

THE eggs, from eight to twelve in number, are  
clear ivory-white in colour.



## TEAL.

## COMMON TEAL.

## PLATE CXCII.—FIGURE I.

*Anas crecca*,  
*Querquedula crecca*,


PENNANT. MONTAGU.  
SELBY.

OF these birds also, the greater number retire to northern regions, such as Iceland, to build, but others remain with us. They have been found on Strensall Common near York.

The nest, which is usually built by the margin of an inland lake, but sometimes near the sea-shore, and in clefts of rocks and stony places, is placed among, and constructed of, heath or grass, or other vegetable substances in moorland and marshy districts, in rushy or boggy places. There is a thick lining of down and feathers: the whole is of rather large size.

The eggs are from ten or eleven to fifteen in number. They have been found under a furze bush without any nest.





TEAL.  
WIGEON.

CXCII

UNIV. OF  
CALIFORNIA







## WIGEON.

## COMMON WIGEON.

## PLATE CXCII.—FIGURE II.

*Anas penelope*,  
*Mareca penelope*,

PENNANT. MONTAGU.  
SELBY. JENYNS.

THESE birds visit northern countries for the most part to breed—Norway, Sweden, and others. They begin to pair by about ‘St. Valentine’s Day.’

Mr. Selby and Sir William Jardine have found the nest in Sutherlandshire, on Loch Laigal, Loch Hope, Loch Shin, and Loch Naver, and also on smaller waters near Lairg.

They build among rushes, grass, flags, and reeds, or low bushes where there are such, and use the former materials and leaves in the construction of the nest, the lining being supplied with down from the bird’s own body. The situation chosen is the neighbourhood of some lake or river, and the nest itself is cleverly concealed.

The eggs are from five to eight, or, Meyer says, ten or twelve in number. They are of a fine cream-white colour.

When the hen bird begins to sit, the male retires into seclusion. The incubation of the eggs continues for twenty-four or twenty-five days. The young, as soon almost as hatched, betake themselves to the water.



## AMERICAN WIGEON.

PLATE CXCH<sup>\*</sup>.—FIGURE I.

*Anas Americana,*  
*Mareca Americana,*

WILSON. AUDUBON.  
 STEPHENS.

THE eggs are from six to eight in number, and of a pale buff colour.

## STELLER'S EIDER.

WESTERN EIDER. WESTERN POCHARD. WESTERN DUCK.

PLATE CXCH<sup>\*</sup>.—FIGURE II.


*Anas dispar,*  
*Fuligula dispar,*  
*Polysticta Stelleri,*  
*Somateria dispar,*

TEMMINCK.  
 SELBY. JENYNS.  
 EYTON.

THIS species breeds, it is said, on high and steep rocks.

The egg is of a pale dusky green colour all over.





UNIV. OF  
CALIFORNIA

AMERICAN WIGEON.  
STELLER'S EIDER.

CXCII\*








48075



PLATE  
CXCIII



EIDER DUCK.  
KING DUCK.

CXCIII



## EIDER DUCK.

ST. CUTHBERT'S DUCK.

PLATE CXCIH.—FIGURE I.

*Anas mollissima,*

PENNANT. MONTAGU.

*Somateria mollissima,*

FLEMING. SELBY.

THE down with which the nests of these birds are lined is made an article of commerce, and in the countries where they are plentiful the people rob the nests two or three times in the season. The eggs are also taken at the same time, so as to make the birds lay again, 'but it is generally found that if they are robbed more than twice, they begin to desert the place, and if pillaged oftener they quit it entirely.'

The Eider Duck begins to pair in March, and to lay about the beginning of the last week in May, that is to say, in this country. Incubation lasts from three weeks to a month. The young, as soon as hatched, are led by the dam to the water, and in some instances it appears certain that they must be carried by her in her bill. The male bird at this period leaves his mate, and repairs to the open sea.

In England the Eiders build so far south as the Fern Islands and Coquet Island. In one of the former the remains of an old lighthouse appears to be a



favourite resort, and within the ruined and deserted walls several have been accustomed to lay. They seem to prefer islands for the purpose, but also to make use of hollows in rocks on the mainland, and sand-banks along the sea-shore. Great numbers are frequently placed close together, so that it is hardly possible to walk among them without treading on some.

It seems that at first the nest has only grass, lichens, or sea-weed, for its composition, but when all, or nearly all, the eggs are laid, down, plucked by the female from herself, is daily added, the quantity being then successively increased. If the bird leaves her charge for a time, the eggs are carefully covered over and concealed, either with down, or grass and leaves, according as each has been supplied.

The eggs, of a light green colour, are usually five in number, but not very unfrequently as many as eight or even ten are found together, two birds having laid in the same place. Both, it appears in that case, sit together in an amicable manner.



## KING DUCK.

KING EIDER.

PLATE CXCIH.—FIGURE II.

*Anas spectabilis*,  
*Somateria spectabilis*,

PENNANT. MONTAGU.  
FLEMING. SELBY.

THE nest of the King Duck, placed on rocks near the sea, is lined with the down of the female, and composed externally of sea-weeds, grass, or mosses.

The eggs are of a pale green colour. They appear to be from four to six in number. The male bird leaves the care of the eggs to the hen, and repairs to the sea.



## VELVET SCOTER.

PLATE CXCIV.—FIGURE II.

*Anas fusca*,  
*Oidemia fusca*,

PENNANT. MONTAGU.  
 FLEMING. SELBY.


A UDUBON writes of this species, that they 'begin to form their nests from the 1st. to the 10th. of June. The nests are placed within a few feet of the borders of small lakes, a mile or two distant from the sea, and usually under the low boughs of bushes, of the twigs of which, with mosses and various plants matted together, they are formed. They are large and almost flat, several inches thick, with some feathers of the female, but no down.'

The eggs are usually six, but sometimes eight or ten in number, of a uniform pale cream-colour, tinged with green.

The males leave the females after incubation has commenced.

A pair had bred on the same water for six or seven years in succession. The young did not quit the pond until they were able to fly; as soon as that is the case the mother bird escorts them to the sea.





UNIV. OF  
CALIFORNIA

SCOTER.  
VELVET SCOTER.

CXCIV







## SCOTER.

COMMON SCOTER. BLACK SCOTER.

PLATE CXCIV.—FIGURE I.

*Anas nigra*,  
*Oidemia nigra*,PENNANT. MONTAGU.  
FLEMING. SELBY.

THE nest of this Duck is described as being placed by the side of water in the most unfrequented situations, often far inland, among brushwood, or on stony ground; and made of grasses, twigs, and the dry leaves and stalks of plants, under the shelter of a willow or birch, or in a dry tuft of herbage, the down of the bird itself being made use of for lining.

After the female has laid, the male birds collect together in flocks, and repair to the coast.

The eggs are from five or six, to seven or eight, or nine, in number;—Thieneman says as many as ten—of a pale buff colour, tinged with green.



## SURF SCOTER.

PLATE CXCIV.—FIGURE II.

*Anas perspicillata*,  
*Oidemia perspicillata*,

TEMMINCK.  
FLEMING. SELBY.

THE nest of this species, placed on the coasts of inland seas, and by the borders of lakes, is formed of dry and withered weeds, flags, and grasses, in the midst of a hummock or tuft of grass. The down of the bird itself serves for the lining. The hollow of the interior is said by Audubon to be six inches across, by two and a half in depth.

The eggs are four, five, or six in number, and of an oval shape; their colour pale yellowish or cream white.





UNIV. OF  
CALIFORNIA

RED-CRESTED WHISTLING DUCK.  
SURF SCOTER.

CXCV







RED-CRESTED WHISTLING DUCK.

RED-CRESTED DUCK. RED-CRESTED POCHARD.

PLATE CXCv.—FIGURE I.

*Anas rufina*,  
*Fuligula rufina*,  
*Mergoides rufina*,

TEMMINCK.  
GOULD. SELBY.  
EYTON.

THE nest of this bird may be termed a bed of rushes.

The eggs are stated to be six or seven in number, and of an uniform olive brown colour.

‘The young are able to fly about the beginning of July; the female is much attached to her young, but the male leaves the nursery as soon as the female begins to sit.’

The figure on the plate is taken from a specimen of the egg of this bird, obligingly forwarded for the use of the present work, by the Rev. H. B. Tristram, Perpetual Curate of Castle Eden, Durham.



## POCHARD.

RED-HEADED POCHARD. COMMON POCHARD.  
POCHARD DUCK. DUN BIRD.

PLATE CXCVI.—FIGURE II.

*Anas ferina*,  
*Fuligula ferina*,  
*Nyroca ferina*,

PENNANT. MONTAGU.  
SELBY. JENYNS.  
FLEMING.


MEYER says, 'about the end of April the Red-headed Pochards pair; there are no quarrels between the male birds, for the female is said to choose her own mate, and the pair remain inseparable, and do not quit the spot until the young brood are hatched and ready to follow them towards the larger lakes, etc., where they consider them out of danger.'

The nest of the Pochard is made among rushes or other coarse herbage, and is lined with feathers. Many nests are placed near each other, in suitable localities, such as osier beds or grassy places.

The eggs are from eight or ten to twelve or thirteen in number, and of a buff white colour.

The Pochard bred, in the year 1855, in the Gardens of the Zoological Society, as Frederick Bond, Esq. has written me word.





UNIV. OF  
CALIFORNIA

FERRUGINOUS DUCK.  
POCHARD.

CXCVI







## FERRUGINOUS DUCK.

WHITE-EYED DUCK. RED DUCK. CASTANEUS DUCK.

PLATE CXCVI.—FIGURE I.

*Anas ferruginea*,

PENNANT. MONTAGU.

*Fuligula nyroca*,

SELBY. JENYNS.

*Nyroca leucophthalma*,

FLEMING.

THIS Duck, like the others of its kindred, builds its nest near rivers, ponds, and marshes. The male bird leaves the female soon after she has begun to sit. The nest is well supplied with down from the breast of the parent mother, as a lining.

The eggs of this species are white, with a slight tinge of green, and nine or ten in number. They are laid by the beginning of June, and are hatched in twenty-two or twenty-three days.

As Frederick Bond, Esq. has informed me, the Ferruginous Duck bred in the year 1854, in the Gardens of the Zoological Society, London; and in that, or the preceding year, also with the Tufted Duck.



## SCAUP.

SCAUP DUCK. SCAUP POCHARD.

PLATE CXCVII.—FIGURE II.


*Anas marila*,  
*Fuligula marila*,  
*Nyroca marila*,

PENNANT. MONTAGU.  
SELBY. JENYNS.  
FLEMING.

THE Scaup builds among the brushwood or other vegetation that is found in swampy grounds, or by lakes, or in stony places near these. Very little nest is formed, the materials, such as they are, being dry grasses, stalks, and leaves, but the eggs are well covered with down. They are, it is said, from five or six to eight or ten in number, of a dull yellowish-brown colour.

The bird begins to lay in May.





UNIV. OF  
CALIFORNIA

TUFTED DUCK.  
SCAUP.

CXCVII







## TUFTED DUCK.

TUFTED POCHARD.

PLATE CXCVII.—FIGURE I.

*Anas fuligula*,  
*Fuligula cristata*,  
*Nyroca fuligula*,

PENNANT. MONTAGU.  
 SELBY. JENYNS.  
 FLEMING.

THESE birds breed along the stony shores of the sides of inland waters, among the cover of vegetation, more less thick, with which they are usually bordered. It is not made till the end of May or beginning of June.

The receptacle for the eggs, for it can hardly be called a nest, is composed of stalks and grasses.

The eggs vary in number from eight to ten. They are of a pale buff colour, with a tinge of green.

The male bird leaves the female after she has begun to sit.

This species, as Frederick Bond, Esq. has informed me, paired with the Ferruginous Duck, as I have mentioned in the account of the latter, in the year 1853 or 1854, in the Gardens of the Zoological Society.



## LONG-TAILED DUCK.

NORTHERN HARELD. CALLOO. COAL-AND-CANDLE-LIGHT.  
 LONG-TAILED SHIELDRAKE. SHARP-TAILED DUCK.

PLATE CXCVIII.—FIGURE I.

*Anas glacialis*,  
*Clangula glacialis*,

PENNANT. MONTAGU.  
 FLEMING. SELBY.


THIS species nidificates about inland lakes, ponds, and bogs, among the plants and low brushwood that afford cover near the water's edge.

The nest is nothing more than a few stems of reeds, rushes, or grass, lined with the down of the mother bird, the quantity being added to as the number of eggs increases. Many breed in the same vicinity.

'About the latter end of April,' says Meyer, 'the birds arrive in pairs at their breeding-places, and in May the female lays. By the end of June or beginning of July the young make their appearance, and are soon led to the sea-shore, and instructed to swim, dive, and provide for themselves. The male, who leaves the care of the nursery to his mate, joins his family on their arriving at that locality, and accompanies them in their migratory journey and winter sojourn.'

The eggs are usually from six to ten in number,





UNIV. OF  
CALIFORNIA

LONG-TAILED DUCK.  
HARLEQUIN DUCK.

CXCVIII







HARLEQUIN DUCK.

101

but twelve have been found. They are of a pale greenish or bluish white colour, tinted with buff.

---

HARLEQUIN DUCK.

HARLEQUIN GARROT.

PLATE CXCVIII.—FIGURE II.

*Anas histrionica*,

PENNANT. MONTAGU.

*Clangula histrionica*,

FLEMING. SELBY.

THE Harlequin builds among low bushes and plants, in the precincts of streams and lakes, in the neighbourhood of the sea, not far from the water's edge; and 'feathers its nest' with down, the outer materials being dry leaves, stalks, grass, and reeds, arranged in a circular manner to the height of two or three inches.

The eggs are from five or six or seven in number, rarely more than the former according to Audubon, but some say as many as eight, ten, or twelve, and of a white or very pale buff colour, with a tinge of green. It is stated that 'the texture of the egg is very fine, but without polish.'



NO. 1111  
 ANGELO

## GOLDEN-EYE.

COMMON GOLDEN-EYE. GARROT. RATTLE-WINGS.

GOLDEN-EYE DUCK.

PLATE CXCIX.—FIGURE II.

<i>Anas clangula</i> ,	PENNANT. MONTAGU.
<i>Anas glaucion</i> ,	BEWICK.
<i>Clangula vulgaris</i> ,	FLEMING.
<i>Clangula chrysophthalmus</i> ,	JENYNS.

THE Golden-Eye builds in the vicinity of lakes and rivers, giving a preference to the latter, particularly such as flow over falls and rapids. The Laplanders place boxes with holes in them in the trees in these localities for the birds to build in, and thus procure the eggs, for the cotes are sure to be resorted to for the purpose of laying in. The nest is made of rushes and other herbage, lined with down. Mr. Hewitson found one in a hole in a tree, ten or twelve feet from the ground.

The eggs are of a greenish hue, and from ten to fourteen in number.





UNIV. OF  
CALIFORNIA

SMEW.  
GOLDEN-EYE.

CXCIX







## SMEW.

LOUGH DIVER. WHITE NUN. WHITE MERGANSER.  
WHITE-HEADED GOOSANDER.

PLATE CXCIX.—FIGURE I.

*Mergus albellus,*

PENNANT. MONTAGU.

THE nest of the Smew is made of dry grass, and lined with the down of the bird itself. It is placed on the ground, upon the banks of lakes and rivers, not far from the water, or in a hollow in a tree.

The eggs are said to be eight or ten, or from that to fourteen in number, and of a yellowish-white colour.



## HOODED MERGANSER.

PLATE CC.—FIGURE I.

*Mergus cucullatus,*

SELBY. JENYNS. EYTON.


*Mergus fuscus,*

LATHAM. GOULD. TEMMINCK.

THE nest, placed in trees, or, when these are wanting, in holes and hollows, in either case by the margins of lakes and rivers, is formed of grass and other herbage, lined with feathers, and down from the breast of the mother bird.

The eggs, eight or ten in number, and of a yellowish-white colour, are laid in May, and hatched in June.





UNIV. OF  
CALIFORNIA

HOODED MERGANSER.  
RED-BREASTED MERGANSER.  
GOOSANDER.







RED-BREASTED MERGANSER.

RED-BREASTED GOOSANDER.

PLATE CC.—FIGURE II.

*Mergus serrator*,

PENNANT. MONTAGU.

THE places chosen by this species for nesting are the vicinage of the sea, and the neighbourhood of lakes and rivers, among reeds and rushes.

These birds build, it seems, on the borders of, and small islands in, lakes, whether of fresh or salt-water, and rivers, preferring such as have a growth of wood, the nest being placed a few yards from the edge, at the foot of a tree, or under the shelter of brushwood, in the midst of fern, grass, nettles, or other wild vegetation. Also in divers other situations, among stones, in a hollow, on the bare ground, at the top of a tall tree, or in the deserted nest of some other bird, or the end of a deep recess. It has been known, moreover, in a bleak and unsheltered situation, on an island in the sea, at some distance from the mainland. The materials of its composition are moss, flags, grass, small roots, and feathers, placed carelessly together, and intermixed with the down of the bird, added to, it appears, as incubation advances.

The eggs are from six or seven, to nine, ten, or



eleven, in number, and of a rich reddish yellow, or brownish fawn-colour. As soon as the females begin to sit, the males quit them for the season. The species appears to be late in its nidification, scarcely beginning to build before the end of May, or the early part of June. The bird sits very close, and will almost allow herself to be trodden on before she will leave the nest.

---

## GOOSANDER.

GREEN-HEADED GOOSANDER. DUN DIVER.

PLATE CC.—FIGURE III.

*Mergus merganser,*

PENNANT. MONTAGU.

*Mergus castor,*

MONTAGU. BEWICK.

THE Goosander builds on small islands in fresh-water lochs in the neighbourhood of the sea, and near the water's edge.

The nest, which is placed under the cover of bushes, in long grass, among stones, or in the hollow of the stump of a decayed tree, is large altogether, being raised to a height of seven or eight inches, on a mass of dead weeds, but the inner and more finished part is only about seven inches and a half across, and four in depth. It is composed of dry grass and small



roots, rather neatly twined together, and lined with the down of the bird.

The eggs, of a long oval shape, are from four to six, seven or eight, and, Yarrell says, ten or fourteen, in number, or more, though rarely; if removed from the nest, as many as thirty have been successively laid: their colour a uniform buff-white or cream-yellow. The figure on the plate is from a pale green variety.

The female has been seen to carry her young on her back, in the same way as described of the Swan. She leads them to the water as soon as they are hatched, or carries them thither in her bill if the nest has been in the hollow of a tree, and at once commences their education.

The males appear to leave the females when the latter have begun to sit, and do not rejoin them till the summer is over: while absent they are only accompanied by one or two females, if any.



## GREAT CRESTED GREBE.

TIPPET GREBE. CRESTED GREBE. GREATER LOON.  
LOON. GAUNT.

PLATE CCI.—FIGURE I.

*Podiceps cristatus*,  
*Colymbus cornutus*,  
*Colymbus cristatus*,

PENNANT. MONTAGU.  
BRISSON.  
LINNÆUS.

THE Loon breeds in fresh water, and makes its nest, such as it is, early in the year—in the month of April. It is a very large and careless mass of weeds, flags, and other water-plants, partly sunk under, and partly raised above the water, the top being slightly hollowed. The general width is about a foot or a little over, the height about half as much.

The eggs are three or four, and occasionally five. Four appears to be the usual average number, but one of them is generally addled. They are concealed by fragments of rushes placed over them, and if these be removed others are added. Their colour is white or greenish white.

Yarrell says, 'The parent birds are very careful of their young, taking them down with them for security under their wings when they dive.' According to Meyer the birds pair for life, and haunt the same nesting-place





UNIV. OF  
CALIFORNIA

GREAT CRESTED GREBE.  
RED-NECKED GREBE.

CCI







year after year, both assisting in the work of nidification. The young swim about as soon as hatched.

---

## RED-NECKED GREBE.

PLATE CCI.—FIGURE II.

*Podiceps rubricollis*,

PENNANT. MONTAGU.

THE nest is made among reeds, flags, and rushes, on the borders of fresh-water lakes and ponds, and is composed of the said materials. It is placed upon the water, and attached to the growing vegetation.

The eggs of the Red-necked Grebe are either white, or white with a slight tinge of blue or green, clouded, Selby says, with deep brown. They are four or five in number.

Several pairs of these birds breed in the same locality.

The male and female both assist in the construction of the nest, and shew much affection for their charge. The young take to the water as soon as they are hatched.



## DUSKY GREBE.

SCLAVONIAN GREBE. HORNED GREBE.

HORNED DABCHICK. BLACK AND WHITE DABCHICK.

PLATE CCII.—FIGURE II.

<i>Podiceps cornutus,</i>	LATHAM. SHAW.
<i>Podiceps obscurus,</i>	LATHAM.
<i>Podiceps Caspicus,</i>	LATHAM.
<i>Colymbus cornutus,</i>	GMELIN.
<i>Colymbus obscurus,</i>	GMELIN.
<i>Colymbus Caspicus,</i>	GMELIN.

THE nest of this bird is large—a mass of reeds, sedges, and other water-plants, placed by the edge of the liquid element, or floated on the surface among the rushes or other such vegetation that there has its natural growth.

The eggs appear to be from two to four or five in number. They are of a white colour, with a faint tinge of blue.





DABCHICK.  
DUSKY GREBE.  
EARED GREBE.

UNIV. OF  
CALIFORNIA

CCII







## EARED GREBE.

EARED DABCHICK.

PLATE CCII.—FIGURE III.

*Podiceps auritus*,  
*Colymbus auritus*,

LATHAM.  
LINNÆUS.

THE bird before us builds its nest of and among reeds and rushes, and places it by the edge of a pond or pool.

The eggs are of a pure white colour, sometimes with and sometimes without a tinge of blue; the surface is rough. They are of a regular oval shape, widest in the middle, and tapering towards each end. Their number, three or four, or, Selby says, four or five.

Incubation is said by Meyer to last three weeks.



## DABCHICK.

LITTLE GREBE. BLACK-CHIN GREBE. DIDAPPER.  
SMALL DOUCKER. LOON.

PLATE CCII.—FIGURE I.

<i>Podiceps minor</i> ,	STEPHENS. FLEMING.
<i>Podiceps Hebridicus</i> ,	LATHAM.
<i>Colymbus Hebridicus</i> ,	GMELIN.
<i>Colymbus minor</i> ,	GMELIN.
<i>Colymbus fluviatilis</i> ,	BRISSON.

THE nest of the Dabchick, which is placed at a little distance from the water, on or among any plants that grow near the sides of rivers, lakes, and ponds, is composed of short pieces of roots, reeds, rushes, and flags, and a considerable quantity of these is occasionally put together, sometimes to the height of a foot or more; when dry the whole naturally becomes very brittle.

The eggs are four, five, or six in number, oval in shape, tapering towards each end, and dull white in colour. The bird is in the habit of covering them over with weeds when leaving the nest for a time, at least, as soon as they are all laid, and it would further appear that the covering is not then removed, but that she sits on both it and the eggs; they become



a good deal stained in consequence. The period of breeding is about the middle of May.

The young take to the water immediately on being hatched.



## GREAT NORTHERN DIVER.

NORTHERN DIVER. GREATEST SPECKLED DIVER. LOON.  
GREAT DOUCKER. IMBER DIVER. IMMER.

PLATE CCIII.

*Colymbus glacialis*,  
*Colymbus Immer*,


LINNÆUS.  
LINNÆUS.

THE nest of this fine bird is placed close to the water's edge, so as to admit of an immediate retreat, if necessary, to that element.

The eggs are two, or sometimes, according to Audubon, three in number. Their colour is a dark olive greenish brown, with a few spots of a darker shade, or purple reddish.

Both birds take their turns in sitting, and when the young are hatched, continue an equal and watchful care over them as long as it is required.





UNIV. OF  
CALIFORNIA

GREAT NORTHERN DIVER.

CCIII









3433





BLACK-THROATED DIVER.  
RED-THROATED DIVER.



## BLACK-THROATED DIVER.

LESSER IMBER. NORTHERN DOUCKER. SPECKLED LOON.

PLATE CCIV.—FIGURE I.

*Colymbus arcticus*,

LINNÆUS. GMELIN.

THIS Diver is found in Europe, in the Lapland and Norwegian seas, and inland waters, even those that exist at a considerable elevation.

In this case also the nest is naturally placed close by the brink of the water, or some small flat island in a lake. It is made of a few straws or stems of plants, placed in a hollow formed or excavated by the bird for the purpose.

The eggs of the Black-throated Diver, usually two in number, are laid on the bare earth, two or three yards from the water's edge. They are of a dark yellowish olive-brown colour, with darker or blackish spots, and of a long and regular oval form.



## RED-THROATED DIVER.

RAIN GOOSE. SPRAT LOON. SPECKLED DIVER.  
 FIRST SPECKLED DIVER. STRIPED DIVER.

PLATE CCIV.—FIGURE II.

<i>Colymbus septentrionalis</i> ,	LINNÆUS. GMELIN.
<i>Colymbus striatus</i> ,	GMELIN.
<i>Colymbus borealis</i> ,	LATHAM.
<i>Colymbus stellatus</i> ,	GMELIN. LATHAM.
<i>Mergus minor</i> ,	BRISSON.

THE situations chosen by this species for laying, are by small pools on low islands in the sea, the margins of inland lakes, and islets in them, and watery places on higher grounds.

The nest is nothing more than a few rushes or blades of grass and moss, mixed, Selby says, with the down of the bird. It is placed among the stones or gravel close to the water's edge, so that if need be, the bird can slide at once into its all but native, and at all times natural and congenial, element.


The eggs are two in number, and equally rounded at both ends. They are of a dull brownish-green colour, rather thickly spotted with dark brown.







UNIV. OF  
CALIFORNIA



GUILLEMOT.

CCV







CCVI



GUILLEMOT.

CCVI



GUILLEMOT.

COMMON GUILLEMOT. FOOLISH GUILLEMOT. TARROCK.  
SCOUT. WILLOCK. LESSER GUILLEMOT. SEA HEN.

PLATES CCV AND CCVI.

*Uria troile,*  
*Uria minor,*

LATHAM. STEPHENS.  
STEPHENS.

THE Guillemot makes no nest, but lays her single egg upon the barren rock. Countless numbers of these birds breed together on the rocks or cliffs that abut upon the ocean, thinking there to find that security, which indeed they would find were it not for the superiority of mind over instinct. Incubation continues for a month. The hen alone sits, and the male does not feed her, but if she be killed after the young are hatched, he continues to bring them up. The old bird is believed to convey her young down to the sea on her back in the evening and when the tide is high, into which it falls or is thrown off by the parent. She appears also to hold it by the shoulder of the wing and so carry it down.

The eggs are very large in proportion to the size of the bird, and more than ordinarily narrowed at one end and widened at the other. They vary in an extraordinary manner, and a description of the principal



varieties only would be almost endless, 'adeo sunt multa.' Some are entirely white, others more or less spotted with brown, and others again bluish green, blotted and streaked with dark reddish brown or black. Some are entirely green. 'The shape of the egg, which is very tapering, prevents it from rolling off into the sea; for when moved by the wind, or other circumstances, it only rolls round its own circle, without changing its first immediate situation.' If the first egg be taken a second is laid, and if the second, a third. It is said that each bird knows its own egg, and that if it be moved she will put it back with her feet. She sits standing if so one may say.







PLATE  
CXXVII



BRUNNICH'S GUILLEMOT.  
RINGED GUILLEMOT.



## BRUNNICH'S GUILLEMOT.

THICK-BILLED GUILLEMOT.

PLATE CCVII.—FIGURE I.

*Uria Brunnichii*,

FLEMING. JENYNS.

THE eggs of this species are, like those of the common kind, most variable in colour. The ground tint is greenish blue. They are, Mr. Hewitson observes, less fantastically streaked and blotted than those of the other, but sometimes very regularly and beautifully marked all over with small spots.



## RINGED GUILLEMOT.

BRIDLED GUILLEMOT.

PLATE CCVII.—FIGURE II.

*Uria lachrymans,*

GOULD. TEMMINCK.

THE figure on the plate is from an egg presented to W. R. Fisher, Esq., by Mr. Wollaston, who had it of Mr. Strickland. The latter gentleman states that the bird breeds regularly at Bridlington, whence he obtained this egg, but in smaller numbers, and in places more difficult of access, than those of the Common Guillemot; and he adds that its eggs are invariably white.



1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860



PLATE OF  
CALCULI



BLACK GUILLEMOT.

CCVIII



## BLACK GUILLEMOT.

SPOTTED GUILLEMOT. COMMON SCRABER.

GREENLAND DOVE.

PLATE CCVIII.

*Uria grylle*,  
*Cephus grylle*,PENNANT. MONTAGU.  
FLEMING.

THIS species pairs about the middle of March, and the eggs are laid in the beginning or more usually by the middle of June. They are hatched in twenty-four days. The bird sits very close, so as to be easily taken on the eggs. Two or more couples have been known to lay under one piece of rock.

The bare earth, or rather the bare rock, or a crevice in it, is the only bed sought for by this species for the purpose of nidification. Mr. Hewitson writes as follows:—‘On some of the islands which present a steep precipice to the sea, they make use of holes or crevices in the rocks, in which the eggs are laid at various distances from the mouth of the hole—from one to two feet, which is most usual, to three or four. On other islands less precipitous, it deposits them in cavities under or between fragments of rock and large stones, with which the beach is strewed. In one place several pairs rear their young ones in crannies between the



stones which form the ruins of an old wall, on the top of a single rock at sea, and at an elevation of fifty or sixty feet above its surface. The Black Guillemot resorts annually to the same holes.'

The eggs are two in number, and of a white colour, with a tinge of green, spotted, blotted, and speckled, more or less, with grey, reddish brown, and very dark brown or blackish. Mr. Hewitson has known one of a pink colour.


Both parents attend to the young until able to fly and dive, when they forage for themselves.







W. H. C. C.



LITTLE AUK.  
PUFFIN.

CCIX



ROTCHE.

LITTLE AUK. COMMON ROTCHE.

PLATE CCIX.—FIGURE I.

*Alca alle,*

PENNANT. MONTAGU.

*Mergulus melanoleucos,*

FLEMING. SELBY.

THE shelter of a chance stone on the earth or rock in some precipitous spot, from whence it can readily drop into the water, or take flight, is all that the Little Auk thinks it necessary to seek or provide in the shape of a nest, unless some cavernous hollow present itself, and this, if suitable, will be taken advantage of by several individuals.

The egg of this small sea-bird is of an uniform pale bluish green; some specimens are spotted with rust-colour. It is of an oval shape, and the ends obtuse.

Several of these birds build in company, from fifty to a hundred or so.

‘While one of the parent birds sits on the eggs, the other may be seen close by, perched on a rock or stone. The young do not leave the nest before they are fully fledged.’



## PUFFIN.

SEA-PARROT. COULTER-NEB. COMMON COULTER-NEB.

PLATE CCIX.—FIGURE II.

*Mormon fratercula*,  
*Alca arctica*,  
*Fratercula arctica*,

GOULD.  
 PENNANT. MONTAGU.  
 FLEMING. SELBY.

THE Puffin breeds in precipitous places on the coast and its adjacent islands, seeming to give a preference to such as are covered with a formation of mould. The nest is made both on cliffs and high rocks, and on the short verdure which obtains on any level places on the sides or summit of such, or among stones and boulders. At first indeed, strictly speaking, there is little or no nest, but as incubation advances, a few grasses may be seen commingled with some feathers of the owners of the hard couch. These birds either take possession of a rabbit-burrow, or dig a hole themselves, as a receptacle for their eggs, frequently to the depth of three feet, and often in a curving direction, and with two entrances. The egg is laid at the farther end. Several will sometimes occupy one hole. A natural cranny or fissure in the rock will equally serve the purpose, or a time-worn hole or hollow in an old wall or ruin. They often build in



company with the Razor-bill, Herring Gull, Kittiwake, and Guillemot, but in separate ranks. Mr. Selby adds that when engaged in digging, which operation is generally performed by the males, they are sometimes so intent upon their work as to allow themselves to be taken by the hand; and the same may also be done when the bird is sitting on its egg. The young are hatched after a month's incubation. In about a like period of time they come to their full feathers, and are able to quit their native burrows, and enter on their ocean of life. Meyer says that where both birds have been killed, others have been known to take charge of the egg or young.

Only one egg is laid, and its colour is white, sometimes spotted with pale grey.



## RAZOR-BILL.

PLATE CCX.

*Alca torda,*

PENNANT. MONTAGU.

ANY slight hollow or cranny in a rock of the cliff overhanging the sea, or even the bare unsheltered surface of the rock itself, a preference being given to the most precipitous places, is chosen by the Razor-bill as a deposit for its single egg. This in the month of April. Even here, however, it is exposed to accidents of different kinds, and is not unfrequently thrown down by a high wind, or some other bird, great numbers frequenting, from the like cause of predilection, the same breeding-places, or is broken by a chance stone or mass of earth dislodged from above; such landslips overwhelm also, at times, the bird herself. The young are borne down to the sea on the back of one or other of the parents. They often build in company with the Kittiwake, the Puffin, the Herring Gull, and the Guillemot, but each kind keeping to itself.

The old ones shew much attachment to their young. The latter are able in July to provide for themselves, but the descent to the sea is not always accomplished with safety. It sometimes happens that in throwing themselves down from the edge of the cliff, to which they are led by their parents, and instructed, as it





UNIV. OF  
CALIFORNIA

RAZOR-BILL.

CCX







were, almost, what to do, they fail in clearing every obstacle below, and the force of the fall in such case is fatal. They are also at times the victims of the onslaughts of Falcons. They lay towards the end of May, or the beginning of June.

The egg is subject to almost endless variety. Its prevailing colour is white, blotted and spotted with blackish brown and reddish brown.



## GREAT AUK.

GAIRFOWL.

PLATE CCXI.

<i>Alca impennis,</i>	PENNANT. MONTAGU. BEWICK. SELBY.
<i>Alca impennis,</i>	FLEMING. JENYNS. GOULD. YARRELL.

THE Great Auk hatches its egg on the bare ground, close above the sea-borde, or in the cleft of a rock. This in the month of June.

The egg is yellowish white, streaked and spotted, principally about the larger end, with black, the ground colour being a dull white tinged with yellow.

These eggs may be said to be 'worth their weight in gold;' I know of about sixty guineas having been given for a pair, and others have fetched much more since then.

I have to thank W. F. W. Bird, Esq., for an offer of a drawing from a specimen in the fine collection of the Right Hon. Lord Garvagh, but the present engraving was already executed: I am equally indebted to my obliging friend.





UNIV. OF  
CALIFORNIA

GREAT AUK.

CCXI










1850



CHAMBERLAIN'S



GREEN CORMORANT.  
CORMORANT.  
GANNET.

CCXII



## CORMORANT.

COMMON CORMORANT. CORVORANT.

PLATE CCXII.—FIGURE II.

*Pelecanus carbo*,  
*Carbo cormoranus*,PENNANT. MONTAGU.  
TEMMINCK.

THE Cormorant naturally prefers an elevated situation for its nest, though in default of such it is obliged to put up with a lowly one. It is well for those of a higher rank in creation than the bird when they can thus readily accommodate themselves to the circumstances in which they are placed. Many pairs congregate together.

In the former case the tops of lofty cliffs are built on, or, as the next best, high trees. Failing these, a bed of rushes is made to serve the purpose on the mainland; or the top of a low island rock is resorted to.

The nest, which is large, is composed of sticks or sea-weeds, heaped up to the height of a couple of feet. The finer portions or grass forming the interior.

The eggs, small in proportion to the size of the bird, are of a pale bluish or greenish white colour, without polish, and of an oblong shape. Three, four, or five are usually laid, but sometimes six. This in



the month of March, April, or May: they are hatched in four weeks.

As soon as the young are able to fend and forage for themselves, which is as soon as they can fly, they are conducted to the sea by their parents, and then left to their own resources.

---

## GREEN CORMORANT.

CRESTED CORMORANT. SHAG. CRESTED SHAG.  
GREEN-CRESTED CORMORANT. CORVORANT.

PLATE CCXII.—FIGURE I.

<i>Pelecanus cristatus</i> ,	PENNANT.
<i>Pelecanus graculus</i> ,	MONTAGU. BEWICK.
<i>Phalacrocorax cristatus</i> ,	SELBY. JENYNS. GOULD.
<i>Carbo cristatus</i> ,	TEMMINCK.

THESE Cormorants make their nests on the ledges and shelves of cliffs or caves over the sea, lower down than the other species. Many pairs, a score or more, frequent the same place. They return to it annually for the breeding-season. Montagu mentions his having counted thirty together on a small rock. The bird sits upright on the eggs.

The nest is a mass of sea-weed, softened off internally with the finer sorts of grass.



The eggs are three, four, or five in number, and their original colour is white, but they soon become stained and discoloured.

---

## GANNET.

COMMON GANNET. SOLAN GANNET. SOLAND GOOSE.  
SOLAN GOOSE.

PLATE CCXII.—FIGURE III.

*Pelecanus bassanus*,  
*Sula alba*,  
*Sula bassana*,

PENNANT. MONTAGU.  
FLEMING.  
BRISSON. SELBY.

GREAT numbers of these birds build together in the same situations, on the sides of precipitous cliffs and rocks, the nests being placed almost close to each other. These are made of sea-weed and grass.

The egg, for there is only one, is white, with a pale, or, I should rather say, a very pale, tinge of blue.

Both parents sit; the period of incubation being about six weeks.



## CASPIAN TERN.

PLATE CCXIII.—FIGURE II.

*Sterna Caspia*, SELBY. JENYNS. EYTON. GOULD.

THE nest is a mere hollow scratched in the sand. The eggs are two or three in number, of a yellowish stone or pale olive green colour, spotted with grey and reddish brown or blackish brown. They are hatched in about twenty days. 'Although the birds use great exertions to prevent an intruder from approaching the nest and eggs, by flying over his head, and making a considerable noise, yet it has been remarked that when they are once disturbed, they do not easily return to their nest, and are said even not to revisit the same spot the following year if they are fired at.' Several hundreds of pairs build together.





UNIV. OF  
CALIFORNIA

CASPIAN TERN.  
SANDWICH TERN.

CCXIII







## SANDWICH TERN.

PLATE CCXIII.—FIGURE I.

*Sterna cantiaca,**Sterna cantiaca,**Sterna Boysii,*

MONTAGU. BEWICK.

JENYNS. GOULD.

FLEMING. SELBY.

THESE birds live together in large colonies, hundreds, and even thousands of pairs crowding together in the same places, the eggs in consequence being so close together that it is difficult to avoid treading on them. The situation chosen is a sandy place, covered at the most with short and bare vegetation. If the nesting-place is approached, the birds fly about in a cloud. They are late in breeding, seldom commencing till the month of June.

The eggs are usually two, but sometimes three or even four in number. Meyer says, 'The bird sits on them during the whole night, but only occasionally during the day, and, as in the preceding species, some few birds remain about the breeding-places, to keep watch during the absence of the rest. It has been asserted that these birds, although laying two or three eggs only for a brood, will, when the eggs are taken out of the nest daily, continue laying for a fortnight.' The eggs vary exceedingly, and are extremely beautiful. They are of a pale yellowish stone-colour, thickly



spotted and marked with deep reddish brown, orange brown, blackish brown, and grey. Some are of a whitish, and others of a dull green ground colour, with spots of a darker shade.







UNIV. OF  
CALIFORNIA



ROSEATE TERN.

CCXIV



## ROSEATE TERN.

PLATE CCXIV.

<i>Sterna Dougallii</i> ,	MONTAGU. BEWICK. FLEMING.
<i>Sterna Dougallii</i> ,	SELBY. JENYNS. GOULD.

THIS species makes its nest among the herbage there may be on low banks of sand or shingle, or upon the bare ground itself.

The eggs are two or three in number; the ground colour yellowish cream white, pale brown, or yellowish olive green, spotted and speckled with grey and brown.



## COMMON TERN.

PLATE CCXV.

<i>Sterna hirundo</i> ,	PENNANT. MONTAGU. BEWICK.
<i>Sterna hirundo</i> ,	FLEMING. SELBY. JENYNS. GOULD.

THE Common Tern lays its eggs on sand or shingle, making little or no nest beyond scooping out a slight hollow: what there is is placed either by the sea-side, or in marshes, on islands, or by the sides of lakes and rivers.

The eggs are variously of a pale blue, pale yellow, green, brown, white, or light dull yellowish or stone-colour, blotted and spotted with grey, dark reddish brown, and blackish brown: three in number. They are laid from the latter end of May to the beginning of July.

The male bird assists the female in the task of sitting during the day, she taking charge of the clutch at night; in fine weather, however, the heat of the sun seems to be thought sufficient warmth, and the bird leaves them to its rays. Ten or twelve pairs of these birds breed together. The young come forth in fifteen or sixteen days, and are able to fly when about three weeks old. The old ones display much anxiety for their safety, and are very clamorous when any one approaches their station, flying round, and frequently even striking against him.





UNIV. OF  
CALIFORNIA

COMMON TERN.

CCXV













PLATE  
CCXVI



ARCTIC TERN.  
SOOTY TERN.

CCXVI



## ARCTIC TERN.

PLATE CCXVI.—FIGURES I, II.

*Sterna arctica*,

FLEMING. SELBY. JENYNS.

*Sterna arctica*,

EYTON. GOULD.

THIS Tern contents itself with a slight hollow scratched out either in the bare sand or grass, a little of the latter forming a lining, by the sea-shore, the borders of islets, or the mouths of rivers. Great numbers build together, and the nests are so closely contiguous that it is hardly possible to avoid walking upon them.

The eggs are two or three in number, rarely four; they are subject to almost endless variety, the ground colour being variously white, green, dull green, brown, pale blue, and pale yellow, spotted with grey and dark reddish-brown. They are usually laid the beginning of June. The period of incubation is about fifteen or sixteen days.



## SOOTY TERN.

PLATE CCXVI.—FIGURE III.

*Sterna fuliginosa*,  
*Onychoprion fuliginosus*,

LATHAM. WILSON.  
GOULD.

NO nest is formed, but the eggs are laid in the sand under trees near the sea-shore, the birds scratching a hollow with their feet, and frequently fitting themselves into it to see that it will answer the purpose.

The eggs are smooth, and of a pale cream-colour, slightly marked with different shades of pale brown and lighter tints of purple. The female, while sitting, is fed by the male in a tender and affectionate manner.







PLATE  
CCXVII



WHISKERED TERN.  
GULL-BILLED TERN.



WHISKERED TERN.

MOUSTACHE TERN.

PLATE CCXVII.—FIGURE I.

*Sterna leucopareia,*

GOULD. TEMMINCK.

THE figure on the plate is from an egg in Mr. Walter's collection, obtained from Dr. Thieneman.



## GULL-BILLED TERN.

PLATE CCXVII.—FIGURE II.

*Sterna Anglica,*

MONTAGU. BEWICK. FLEMING.

*Sterna Anglica,*

SELBY. JENYNS. EYTON. GOULD.

THE nesting-places of this species are marshy grounds not far from the sea-beach. A small hollow is scratched in the sand or gravel, and in this a few stems of grass and small roots are placed.

The eggs are two, three, or four in number, of a pale, as also of a dull greenish or yellowish olive-colour, marked and spotted with pale grey, reddish brown of two shades, and olive brown. J. R. De Capel Wise, Esq., of Lincoln College, Oxford, has favoured me with the loan of a specimen.







PLATE 18



LESSER TERN.  
BLACK TERN.



## LESSER TERN.

## LITTLE TERN.

## PLATE CCXVIII.—FIGURE I.

*Sterna minuta*,

PENNANT. MONTAGU. BEWICK.

*Sterna minuta*,

FLEMING. SELBY. JENYNS. GOULD.

THE bird before us lays in the sand or gravel along the sea-shore, as also by the borders of rivers: several pairs breed close together.

The eggs are generally two, or sometimes three in number, of a pale brown, dull green, or stone-colour, spotted and speckled with grey and dark chesnut brown. They are laid by the first week in June.

The young are hatched in a fortnight: they are able to fly by about the second week in July. Audubon says that they are fed for a time on the wing by both parents.



## BLACK TERN.

PLATE CCXVIII.—FIGURE II.

*Sterna nigra*,  
*Sterna nigra*,  
*Sterna fissipes*,  
*Sterna nævia*,

FLEMING. SELBY.  
 JENYNS. GOULD.  
 PENNANT.  
 PENNANT.

THESE birds build together in great numbers; they choose for the purpose swampy places among moors or fens. Their nests are either placed in some slight mound of earth, or tuft of grass or rushes, close to, or floated on the water, among the kindred stems and stalks of flags and reeds, and with grass for the inner portion.

The eggs are three in number, and still oftener four, and of a rather dark olive-green or olive-brown colour, blotted and spotted with deep brown or black, principally at the larger end.

The young are produced in fifteen or sixteen days.







PLATE  
CCXIX



WHITE-WINGED BLACK TERN.

CCXIX



WHITE-WINGED BLACK TERN.

SILVER-WINGED BLACK TERN.

PLATE CCXIX.

*Sterna leucoptera*,

MEYER.

THE eggs of this species are three or four in number, with many greyish spots, and some larger blackish-red ones, the ground colour being dull yellowish-olive. They are of a rotund form.

The male and female birds sit on them in turn, and shew much anxiety for their safety, flying at and about all intruders.

The figure on the plate is from an egg forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## NODDY.

BLACK NODDY. NODDY TERN.

PLATE CCXX.—FIGURE I.

*Sterna stolidus*,  
*Anous stolidus*,


JENYNS. GOULD.  
EYTON.

THE nesting-places of this bird are banks and rocks.

The nest, which is placed in low trees or bushes, is made of twigs and dry grass, and that of the former year is often repaired, being thus raised to the height of nearly a couple of feet. Thousands of birds build together.

The eggs, three in number, are of a reddish yellow colour, spotted with dull red and purple. They are thought very good to eat.





UNIV. OF  
CALIFORNIA

NODDY.  
SABINE'S GULL.

CCXX







## SABINE'S GULL.

PLATE CCXX.—FIGURE II.

*Larus Sabini*,  
*Larus Sabineus*,  
*Xema Sabini*,

JENYNS.  
TEMMINCK.  
EYTON. GOULD.

THIS bird lays its eggs on the bare earth.

They are two or three in number, olive brown in their ground colour, numerous spotted with darker brown. They are of a regular shape, and not much pointed.

The parent birds, when engaged with their young, exhibit much anxiety for them, and dash about intruders in a perturbed manner.

The young are hatched in the latter end of July.

The figure on the plate is from an egg forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



## LITTLE GULL.

PLATE CCXXI.—FIGURE I.

<i>Larus minutus,</i>	MONTAGU. BEWICK.
<i>Larus atricilloides,</i>	GMELIN. FLEMING.
<i>Chroiocephalus minutus,</i>	EYTON.
<i>Xema minutus,</i>	GOULD.

THE eggs have a ground colour of a greenish ochre, spotted with grey and reddish brown.

---

## BUONAPARTE'S GULL.


BUONAPARTIAN GULL.

PLATE CCXXI.—FIGURE II.

<i>Larus Buonapartii,</i>	THOMPSON. AUDUBON.
---------------------------	--------------------

THE eggs are of a light brown colour, spotted with grey and two shades of brown.





UNIV. OF  
CALIFORNIA

LITTLE GULL.  
BUONAPARTE'S GULL.

CCXXI.








3333



UNIV. OF  
CALIFORNIA



MASKED GULL.  
BLACK-HEADED GULL.



MASKED GULL.

BROWN-HEADED GULL.

PLATE CCXXII.—FIGURES I, II.

<i>Larus capistratus,</i>	FLEMING. JENYNS.
<i>Chroicocephalus capistratus,</i>	EYTON.

THE eggs are described as of a greyish green colour, with dark spots.

The specimens from which the engraving is made were furnished for the use of this work by J. R. De Capel Wise, Esq., of Lincoln college, Oxford.



## BLACK-HEADED GULL.

PEEWIT GULL. RED-LEGGED GULL.

PLATE CCXXII.—FIGURE III.

*Larus ridibundus*,

LINNÆUS. LATHAM.

THIS pretty-looking bird resorts to fenny places and the sides of pools and inland waters and their islands to breed, and vast multitudes congregate together for the purpose, as well both near the sea and far from it, even to the lands adjoining the sea itself, if low and marshy.

The nest is flat, and a composition of grass or the tops of reeds and sedge, placed, perhaps, on a tuft of rushes or other such herbage.

If the first set of eggs be taken a second is laid, and a third if the second, but in such cases they are less each time in size. They are valued as food, and in some places are farmed for the purpose.

The eggs, three or four in number, are laid the middle or end of April, or beginning of May, chiefly at the latter season, and are hatched the end of May or early in June. They vary exceedingly in colour and markings; some are light blue, others yellow, and others green, red, or brown. Some have scarcely any spots, and others are thickly covered with marks of



different shades of brown and reddish brown. One beautiful variety has been described, the ground colour a very light greenish white, blotted with two shades of rich brown. In some instances they are found entirely white.

The young birds leave the nest and betake themselves to the water as soon as hatched.

Sir William Jardine writes, 'They are particular in the choice of a breeding-place, at least some which we would think suited for them are passed or deserted, and others more unlikely are selected. We possess a reedy loch which was for many years a haunt of these birds, but the edges were planted and they left it; ten years afterwards, and when the plantation had grown up, a few pairs returned, and in time increased to a large colony, when an artificial piece of water had been made by damming up a narrow pass in an extensive muir, nearly two miles distant; thither the Gulls resorted the following spring, leaving their ancient ground; and they have been increasing in numbers for some years past.'



MEDITERRANEAN BLACK-HEADED  
GULL.

PLATE CCXXII\*.—FIGURE I.

*Larus melanocephalus*,

NATTERER.

THE nest is made of small twigs.  
The eggs, from two to three in number, are  
of a pale stone-colour.

---

GREAT BLACK-HEADED GULL.



PLATE CCXXII\*.—FIGURE II.

*Larus ichthyætus*,

PALLAS.

THE bare sand serves as a nest.  
The eggs, three in number, are of a dull  
yellowish stone-colour, streaked and blotted with black  
and brown.





UNIV. OF  
CALIFORNIA

MEDITERRANEAN BLACK-HEADED GULL.  
GREAT BLACK-HEADED GULL.

CCXXXII







3533



LAUGHING GULL.  
KITTIWAKE.



LAUGHING GULL.  
KITTIWAKE.

CCXXIII



## LAUGHING GULL.

PLATE CCXXIII.—FIGURE I.

*Larus atricilla*,  
*Xema atricilla*,

MONTAGU. FLEMING.  
GOULD.

THE present species builds in marshy places near the sea-shore. Meyer asserts that during fine weather this bird sits on the eggs only at night, leaving the breeding-place early in the morning for the purpose of feeding; but that, when the weather is wet or cold, she remains on the eggs, and takes care of them also during the day.

The nest is made of dry grass and sea-weeds, arranged to a height of two or three inches, the interior being from four and a half to five inches wide, and one and a half deep.

Audubon mentions a curious instance of two nests being built as it were into one; each pair of birds, both male and female, living in the closest companionship, in both senses of the word. In some instances the sand alone is hollowed into a nest.

The eggs are three in number at the most; their usual colour is brownish olive, spotted and blotted with dull reddish brown, the said markings being most numerous about the larger end. They are considered very good to eat.



The old birds exhibit much care for their charge.  
The young ones quit the nest soon after being hatched.

The plate is from an egg forwarded by Mr. Wise.

---

## KITTIWAKE.

TARROCK, (YOUNG.)

PLATE CCXXIII.—FIGURES II, III.

*Larus tridactylus*,

LATHAM.

*Larus rissa*,

LINNÆUS.

*Rissa Brunnichii*,

STEPHENS.

THE Kittiwake lays its eggs high up on or in any small ledge or cleft at the side of a steep and rocky cliff by the sea. These are often so narrow and apparently insecure, that Selby says the young seem instinctively aware of their perilous situation, whence sometimes the least movement would precipitate them into the waves beneath, and are observed seldom to change their attitude in the nest till sufficiently fledged to be able to provide for their own safety. Immense numbers of these birds build together, so much so as completely to whiten the places where they assemble for the purpose, and to give them the appearance of being covered with snow. They often are found in



company with the Razor-bill, Guillemot, and Puffin.

The nest is a mixture of grass and other dry herbage with sea-weed.

The time for laying is the latter end of May, or the beginning of June.

The eggs are usually two, but occasionally, though very rarely, three in number. They differ much in their colouring and markings; the prevailing tint is stone-colour with a tinge of olive, much spotted with grey and brown of two shades; or greyish-white faintly tinged with brown, and blotted with dark brown and purple grey.



## IVORY GULL.

PLATE CCXXIV.—FIGURE I.

*Larus candidus,*

FLEMING.

*Larus eburneus,*



GMELIN. LATHAM.

THIS Gull builds on cliffs and rocky shores in the extreme north; and the nest is said to be a layer of sea-weed.

The eggs are reported to be two or three in number; they are of a cream-white colour, spotted with grey, reddish brown, and brownish black.

The specimen from which the plate is taken was had by the Rev. H. B. Tristram, Perpetual Curate of Castle Eden, Durham, from the west side of Greenland, in the year 1853.





IVORY GULL.  
ICELAND GULL.

UNIV. OF  
CALIFORNIA

CCXXIV







## ICELAND GULL.

LESSER WHITE-WINGED GULL.

WHITE-WINGED SILVERY GULL.

PLATE CCXXIV.—FIGURE II.

<i>Larus Icelandicus,</i>	EDMONSTONE. SELBY.
<i>Larus argentatus,</i>	SABINE. TEMMINCK.
<i>Larus leucopterus,</i>	BUONAPARTE.
<i>Larus arcticus,</i>	MACGILLIVRAY.
<i>Larus glaucoides,</i>	TEMMINCK.

THE Iceland Gull builds on the face of a precipice or cliff, but at a rather low height. It shares the place with other species.

The eggs are of an olive green colour, spotted with two shades of brown.

The Rev. H. B. Tristram has favoured me also with the egg of this bird to be made use of for the plate. It was procured from the White Sea in 1852.



## COMMON GULL.

SEA MEW. WINTER MEW.

PLATE CCXXV.

*Larus canus,*  
*Larus hybernus,*  
*Larus fuscus,*

LINNÆUS.  
GMELIN.  
RAY.

THE nest of this Gull, which is rather large, is only a slight comminglement of grass if placed on the turf, or a larger assortment of this material and sea-weed if built upon the rock; precipitous places are used as well as the lower ground, and in one instance Sir William Jardine mentions his having known several nests placed on the wall of an old ruin. Both the sea-coasts and the shores of inland lakes and marshy islands are resorted to for nidification.

These birds appear to keep aloof in their building as much as may be from their congeners. Large numbers breed together in many cases, and in others a few pairs, or even a single pair by themselves.

The eggs, two or sometimes three in number, are of various shades of colour—light blue, yellowish white, pale yellow, green-brown, and dark olive brown, spotted and blotted irregularly with deeper brown, grey, and black.





UNIV. OF  
CALIFORNIA

COMMON GULL.

CCXXV













LESSER BLACK-BACKED GULL.  
HERRING GULL.



LESSER BLACK-BACKED GULL.

YELLOW-LEGGED GULL.

PLATE CCXXVI.—FIGURE I.

*Larus fuscus*,

LINNÆUS. GMELIN.

THE present species builds on bare and barren islands, both those of the sea-coast and those of inland lakes, as well as on the margin of the mainland. Steep and precipitous cliffs are used, as well as low marshy moors and sandy places a little way inland.

The nest is a tolerable quantity of grass, with occasionally, though rarely, some sea-weed placed loosely together in any slight hollow in the rock or the adjacent herbage, the latter least frequently.

The first or second week in June is the time of building.

The eggs, two, three, or four in number, are of almost every variety of shade of green and brown, olive green, olive-brown, and stone-colour, thickly spotted with grey, brown of two shades, and brownish-black.



## HERRING GULL.

SILVERY GULL.

PLATE CCXXVI.—FIGURE II.

*Larus argentatus,*

BRUNNICH. GMELIN.

*Larus glaucus,*

TEMMINCK.

*Larus marinus,*

LATHAM.

THE Herring Gull makes a nest of dry grass, plants, and sea-weed on the ledges and small grassy or stony places that occur along the side of a cliff, and towards the top. It is of a considerable size outside, the margin being flat, and the inner part small and round.

Great numbers of these birds breed together, and in close neighbourhood also with other species,—the Kittiwake, Razor-bill, Puffin, and Guillemot; but still distinct.

The eggs are two, or more properly three, in number. They are of a light olive-brown colour, spotted over with darker brown of two shades; some deep olive-green, blotted over with blackish-brown. They are laid about the middle of May, and the young take wing about the 20th. of July.







PLATE 38  
CALIFORNIA



GREAT BLACK-BACKED GULL.  
GLAUCOUS GULL.



GREAT BLACK-BACKED GULL.

GREATER BLACK-BACKED GULL.

GREAT BLACK AND WHITE GULL. WAGEL (YOUNG.) COBB.

PLATE CCXXVII.—FIGURE I.

<i>Larus marinus</i> ,	LINNÆUS. GMELIN. BRUNNICH.
<i>Larus niger</i> ,	BRISSON.
<i>Larus ævius</i> ,	GMELIN.

THE Great Black-backed Gull builds chiefly in marshes and low undrained moors, as also in and on the cliffs and rocky islets of the sea-coast and of inland lakes, making its nursery of dry grass, seaweeds, and sticks. Many resort to the same place. The nest is of large size. Both male and female assist in its construction.

The eggs are three in number. They are considered very good to eat, and great numbers are taken for the purpose, the first and second clutch being removed, and the bird then laying a third time. Their colour is yellowish-brown, with a tinge of green, a little spotted with bluish-grey and dark brown.



## GLAUCOUS GULL.

BURGOMASTER.

PLATE CCXXVII.—FIGURE II.

*Larus glaucus*,

BRUNNICH. GMELIN.

THIS bird is said to build upon the sea-beach, just above high-water mark, as well as on the ledges of steep cliffs, and island rocks in the sea. The nests are composed of dry sticks and sea-weeds, with some lining of grasses, the whole laid together to the depth of about half a foot, the width being about two feet. They are tolerably strong, though inartificially constructed.

The eggs are stated to be of a pale purple-grey colour, with scattered spots of brown, and pale bluish purple.


The plate is from a specimen forwarded by J. R. De Capel Wise, Esq., of Lincoln College, Oxford.



1850



PLATE  
CCXXVIII



POMERINE SKUA.  
SKUA.

CCXXVIII



## SKUA.

COMMON SKUA. SKAU GULL. BROWN GULL. BONXIE.

PLATE CCXXVIII.—FIGURE II.

<i>Cataractes vulgaris</i> ,	FLEMING.
<i>Cataractes skua</i> ,	STEPHENS.
<i>Catharacta skua</i> ,	BRUNNICH.
<i>Larus cataractes</i> ,	LINNÆUS. GMELIN. LATHAM.

THE nest of the Skua is of large size, as well as somewhat carefully constructed; the materials used being grass, lichens, moss, and heath. The bird places it on the tops of the mountains or cliffs in the neighbourhood of the sea, but not on the rocks themselves. They build separately in pairs.

The eggs are only two. They are laid towards the end of June.



## POMERINE SKUA.

POMERINE JAGER.

PLATE CCXXVIII.—FIGURE I.

*Cataractes pomarinus*,

STEPHENS.

*Lestris pomarinus*,

TEMMINCK. SABINE.

THIS Skua Gull builds by the margin of lakes in marshy places, as well as in more rocky ones.

The nest is composed of different grasses and moss.

The eggs are two or three in number, and of a yellowish-grey colour, spotted with blackish brown.


J. R. De Capel Wise, Esq., of Lincoln College, Oxford, obligingly forwarded the egg from which the engraving is taken.







524. 54  
C. 220084



BUFFON'S SKUA.  
RICHARDSON'S SKUA.

CCXXIX



## RICHARDSON'S SKUA.

RICHARDSON'S ARCTIC GULL.

PLATE CCXXIX.—FIGURE II.

*Lestris Richardsonii*,

JENYNS. GOULD.

*Larus parasiticus*,

MONTAGU.

THESE birds build both separately and in companies —twenty or thirty pairs together. They make their nests on some raised part of a marshy place, or the top of a moor waste; the heath, moss, or grass thereon being imbedded into a nest.

The eggs are two in number. They are laid early in June. They are of an olive-brown colour, spotted with dark brown.



## LONG-TAILED SKUA.

BUFFON'S SKUA. ARCTIC SKUA. TEAZER. ARCTIC GULL.  
ARCTIC JAGER. BLACK-TOED GULL.

## PLATE CCXXIX.—FIGURE I.

<i>Cataractes parasiticus</i> ,	FLEMING.
<i>Lestris parasiticus</i> ,	JENYNS. EYTON. GOULD.
<i>Stercorarius cephus</i> ,	SHAW.
<i>Stercorarius longicaudus</i> ,	BRISSON.
<i>Larus parasiticus</i> ,	LINNÆUS.
<i>Larus crepidatus</i> , (young)	GMELIN. LATHAM.
<i>Lestris crepidatus</i> , (young)	TEMMINCK.
<i>Cataracta cephus</i> ,	BRUNNICH. RAY.

THIS Skua Gull breeds on the shores of the sea and those of the larger rivers, as also on barren heaths at some distance inland: many birds build together. The nest is composed of dry grass. The old birds are very daring in defence of their young.


The eggs are in number two, of a pale olive-green colour, with irregular blots of reddish-brown.







CCXXX



FULMAR.

CCXXX



## FULMAR.

FULMAR PETREL. NORTHERN FULMAR. PETREL FULMAR.

PLATE CCXXX.

*Procellaria glacialis*,

PENNANT. MONTAGU. BEWICK.

THE Fulmar builds on the small grassy shelves that occur on the front of high and inaccessible precipices, the result of the dilapidations that time works in even the hardest rock. Numberless pairs of these birds build close together.

A slight hollow in the turf, lined with grass and tufts of the sea-pink, forms the nest.

The single egg is of very large size, white, and of a brittle texture.

The young are hatched about the middle of June, the eggs having been laid the beginning of that month.

The Rev. H. B. Tristram obtained this egg from Iceland, and has obligingly given me the use of it for this work.



## MANX SHEARWATER.

SHEARWATER PETREL. MANX PUFFIN. MANKS PETREL.

PLATE CCXXXI.—FIGURE II.

*Procellaria puffinus*,  
*Procellaria anglorum*,  
*Puffinus anglorum*,

PENNANT. MONTAGU.  
 JENYNS. TEMMINCK.  
 FLEMING. SELBY.

THESE birds resort for the purpose of incubation to the highest grassy parts of small rocky islands and the kindred shores of the mainland, as also to sandy places, where they breed in burrows, going in to a depth of about two feet. The excavating of these appears to occupy a considerable time. Both birds sit, and that so closely as to allow themselves to be taken in the hand.


They appear to be very variable in the time of laying, from the end of May to the end of June.

The nest is at most and best but a slight collection of fern leaves and withered stems of other plants; frequently none is formed, but the sand alone suffices the bird.

Only one egg is produced; it is large in size in proportion to the bulk of the bird, perfectly white, and of an oval form, but both ends obtuse.

This egg was taken by the Rev. H. B. Tristram, in Orkney. The plate is figured from it.





UNIV. OF  
CALIFORNIA

BULWER'S PETREL.  
MANX SHEARWATER.

CCXXXI







BULWER'S PETREL.

PLATE CCXXXI.—FIGURE I.

*Procellaria Bulwerii*,  
*Thalassidroma Bulwerii*,

JARDINE. SELBY.  
GOULD.

THIS egg was obtained from Madeira by the Rev.  
H. B. Tristram, and by him politely lent to me  
for the use of this work.



## WILSON'S PETREL.

PLATE CCXXXII.—FIGURE II.

*Procellaria Wilsoni*,  
*Thalassidroma Wilsoni*,

JENYNS.  
TEMMINCK.

THESE birds appear to build in numbers, about the beginning of June, on sand islands, in which they burrow small holes to the depth of two feet or two and a half: the necessary bedding provided is a little grass.

The egg is a single one, and its colour white.

The young are able to go to sea by the beginning of August.

The egg figured in the plate was lent to me by the Rev. H. B. Tristram, Curate of Castle Eden, Durham. He had it from Newfoundland.





UNIV. OF  
CALIFORNIA

STORM PETREL.  
WILSON'S PETREL.  
FORK-TAILED PETREL.

CCXXXII







## FORK-TAILED PETREL.

LEACH'S PETREL. FORK-TAILED STORM PETREL.

PLATE CCXXXII.—FIGURE III.

*Procellaria Leachii*,

BEWICK. JENYNS.

*Procellaria Bullockii*,

FLEMING.

*Thalassidroma Bullockii*,

EYTON.

*Thalassidroma Leachii*,

GOULD. TEMMINCK.

THIS Petrel breeds in burrows in the sand, among stones, or adopts a hole or hollow in the cleft of a huge wave-worn rock for its nest.

The egg is white—only one is laid. It is of large proportionate size.

I have to thank the Rev. H. B. Tristram for the loan of the egg from which the plate is taken. It was had by him from St. Kilda.



## STORM PETREL.

MOTHER CAREY'S CHICKEN. COMMON STORM PETREL.  
STORMY PETREL.

PLATE CCXXXII.—FIGURE I.

<i>Procellaria pelagica</i> ,	PENNANT. MONTAGU. BEWICK.
<i>Procellaria pelagica</i> ,	FLEMING. JENYNS. TEMMINCK.
<i>Thalassidroma pelagica</i> ,	SELBY. GOULD.

A few pieces of stalks of plants, dried grass, or sea-pinks, with a stray feather or two, are all the nest. The bird sits very close, and will allow herself to be taken sooner than forsake her charge.

The egg is white, and somewhat of an oval shape. It is very frequently surrounded about the base with a ring of faint dull-coloured pink or fine rust-coloured spots.

The Rev. H. B. Tristram took the egg from which the plate is coloured in Orkney.



## INDEX.

A		Vol. Page	
Aberdevine . . .	ii	18	
<i>Accipiter alaudarius</i> . . .	i	35	
“ <i>fringillarius</i> . . .	i	41	
“ <i>palumbarius</i> . . .	i	39	
Accentor, Alpine . . .	ii	59	
<i>Accentor alpinus</i> . . .	ii	59	
“ <i>modularis</i> . . .	ii	60	
<i>Acrocephalus aquaticus</i> . . .	ii	89	
“ <i>palustris</i> . . .	ii	95	
<i>Acrocephalus Bellonii</i> . . .	iii	31	
African Heron . . .	iii	4	
<i>Agrodona campestris</i> . . .	i	152	
<i>Alauda alpestris</i> . . .	i	162	
“ <i>arborea</i> . . .	i	165	
“ <i>arvensis</i> . . .	i	167	
“ <i>brachydactyla</i> . . .	i	164	
“ <i>campestris</i> . . .	i	152	
“ “ . . .	i	154	
“ “ . . .	i	156	
“ “ <i>spinoletta</i> . . .	i	156	
“ “ . . .	i	160	
“ <i>cornuta</i> . . .	i	162	
“ <i>cristata</i> . . .	i	168	
“ <i>cristatella</i> . . .	i	165	
“ <i>grandior</i> . . .	i	152	
“ <i>Ludoviciana</i> . . .	i	156	
“ <i>minor</i> . . .	i	158	
“ <i>mosellana</i> . . .	i	152	
“ <i>nemorosa</i> . . .	i	165	
“ <i>obscura</i> . . .	i	160	
“ <i>pensylvanica</i> . . .	i	156	
“ <i>petrosa</i> . . .	i	160	
“ <i>pratensis</i> . . .	i	154	
“ <i>rufra</i> . . .	i	156	
“ <i>rufa</i> . . .	i	156	
“ <i>spinoletta</i> . . .	i	153	
“ “ . . .	i	156	
“ <i>trivialis</i> . . .	i	154	
“ “ . . .	i	158	
<i>Alauda vulgaris</i> . . .	i	167	
<i>Alca alle</i> . . .	iii	123	
“ <i>arctica</i> . . .	iii	124	
“ <i>impennis</i> . . .	iii	126	
“ <i>torda</i> . . .	iii	127	
<i>Alcedo alcyon</i> . . .	i	92	
“ <i>ispida</i> . . .	i	90	
Alp . . .	ii	26	
Alpine Accentor . . .	ii	59	
Alpine Swift . . .	i	135	
“ Warbler . . .	ii	59	
<i>Aluco flammeus</i> . . .	i	54	
“ <i>minor</i> . . .	i	54	
American Bittern . . .	iii	13	
American Purple Martin . . .	i	138	
American Wigeon . . .	iii	88	
American Yellow-billed Cuckoo . . .	i	126	
<i>Ampelis garrulus</i> . . .	i	113	
<i>Anas acuta</i> . . .	iii	80	
“ <i>Americana</i> . . .	iii	88	
“ <i>anser</i> . . .	iii	63	
“ <i>boschas</i> . . .	iii	81	
“ <i>Canadensis</i> . . .	iii	71	
“ <i>clangula</i> . . .	iii	102	
“ <i>clypeata</i> . . .	iii	78	
“ <i>crecca</i> . . .	iii	86	
“ <i>cygnus</i> . . .	iii	72	
“ <i>dispar</i> . . .	iii	88	
“ <i>Egyptiaca</i> . . .	iii	70	
“ <i>erythropus</i> . . .	iii	68	
“ <i>ferina</i> . . .	iii	96	
“ <i>ferruginea</i> . . .	iii	97	
“ <i>ferus</i> . . .	iii	63	
“ <i>fuligula</i> . . .	iii	99	
“ <i>fusca</i> . . .	iii	92	
“ <i>glacialis</i> . . .	iii	100	
“ <i>glaucion</i> . . .	iii	102	
“ <i>histrionica</i> . . .	iii	101	
“ <i>leucopsis</i> . . .	iii	68	
“ <i>marila</i> . . .	iii	98	
“ <i>mollissima</i> . . .	iii	89	



	Vol. Page		Vol. Page
<i>Anas nigra</i> . . . . .	iii 93	Arctic Jager . . . . .	iii 164
" <i>olor</i> . . . . .	iii 74	" Skua . . . . .	iii 164
" <i>perspicillata</i> . . . . .	iii 94	Arctic Tern . . . . .	iii 137
" <i>querquedula</i> . . . . .	iii 83	<i>Ardea æquinoctialis</i> . . . . .	iii 8
" <i>rufina</i> . . . . .	iii 95	" <i>affinis</i> . . . . .	iii 8
" <i>rutilla</i> . . . . .	iii 77	" <i>alba</i> . . . . .	iii 6
" <i>segetum</i> . . . . .	iii 65	" " <i>major</i> . . . . .	iii 6
" <i>spectabilis</i> . . . . .	iii 91	" <i>Botaurus</i> . . . . .	iii 4
" <i>strepera</i> . . . . .	iii 79	" <i>candida</i> . . . . .	iii 6
" <i>Tadorna</i> . . . . .	iii 76	" <i>Caspica</i> . . . . .	iii 4
Andalusian Hemipode . . . . .	ii 150	" <i>Castanea</i> . . . . .	iii 9
Andalusian Quail . . . . .	ii 150	" <i>ciconia</i> . . . . .	iii 14
Andalusian Turnix . . . . .	ii 150	" <i>cinerea</i> . . . . .	iii 3
<i>Anous stolidus</i> . . . . .	iii 144	" <i>comata</i> . . . . .	iii 9
<i>Anser albifrons</i> . . . . .	iii 67	" <i>cristata</i> . . . . .	iii 3
" <i>bernicle</i> . . . . .	iii 68	" <i>danubialis</i> . . . . .	iii 12
" <i>brachyrhynchus</i> . . . . .	iii 66	" <i>egretta</i> . . . . .	iii 6
" <i>brenta</i> . . . . .	iii 68	" <i>Egretloides</i> . . . . .	iii 6
" <i>Canadensis</i> . . . . .	iii 71	" <i>erythropus</i> , var. <i>B.</i> . . . .	iii 9
" <i>Egyptiacus</i> . . . . .	iii 70	" <i>Gardeni</i> . . . . .	iii 10
" <i>erythropus</i> . . . . .	iii 67	" <i>gazzella</i> . . . . .	iii 7
" <i>ferus</i> . . . . .	iii 63	" <i>grus</i> . . . . .	iii 1
" " . . . . .	iii 65	" <i>lentiginosa</i> . . . . .	iii 13
" <i>leucopsis</i> . . . . .	iii 68	" <i>major</i> . . . . .	iii 3
" <i>palustris</i> . . . . .	iii 63	" <i>marsigli</i> . . . . .	iii 9
" <i>phœnicopus</i> . . . . .	iii 66	" <i>minor</i> . . . . .	iii 13
" <i>ruficollis</i> . . . . .	iii 69	" <i>minuta</i> . . . . .	iii 12
" <i>torquatus</i> . . . . .	iii 68	" <i>Mokoho</i> . . . . .	iii 13
<i>Anthus aquaticus</i> . . . . .	i 153	" <i>nigra</i> . . . . .	iii 15
" " . . . . .	i 156	" <i>Nycticorax</i> . . . . .	iii 10
" " . . . . .	i 160	" <i>obscura</i> . . . . .	iii 10
" <i>arboreus</i> . . . . .	i 158	" <i>pumila</i> . . . . .	iii 9
" <i>campestris</i> . . . . .	i 152	" <i>purpurea</i> . . . . .	iii 4
" " . . . . .	i 160	" <i>Ralloides</i> . . . . .	iii 9
" <i>Ludovicianus</i> . . . . .	i 156	" <i>rußata</i> . . . . .	iii 8
" <i>minor</i> . . . . .	i 158	" <i>Senegalensis</i> . . . . .	iii 9
" <i>montanus</i> . . . . .	i 153	" <i>Squaiotta</i> . . . . .	iii 9
" " . . . . .	i 156	" <i>stellaris</i> . . . . .	iii 11
" <i>paludosa</i> . . . . .	i 152	" " <i>major</i> . . . . .	iii 4
" <i>petrosus</i> . . . . .	i 160	" " <i>var.</i> . . . .	iii 13
" <i>pratensis</i> . . . . .	i 154	" <i>variegata</i> . . . . .	iii 4
" <i>Ricardi</i> . . . . .	i 151	<i>Ardeola minuta</i> . . . . .	iii 12
" <i>rufescens</i> . . . . .	i 152	<i>Arenaria calidris</i> . . . . .	ii 171
" <i>rupestris</i> . . . . .	i 160	" <i>cinerea</i> . . . . .	ii 167
" <i>spinoletta</i> . . . . .	i 153	" <i>vulgaris</i> . . . . .	ii 171
" " . . . . .	i 156	Ash-coloured Sandpiper . . . . .	iii 41
<i>Apterus tridactylus</i> . . . . .	i 125	" Shrike . . . . .	i 59
Aquatic Warbler . . . . .	ii 89	<i>Aster palumbarius</i> . . . . .	i 39
<i>Aquila albicilla</i> . . . . .	i 6	Auk, Great . . . . .	iii 128
" <i>chrysaëtes</i> . . . . .	i 8	Auk, Little . . . . .	iii 123
" <i>haliaëtos</i> . . . . .	i 13	Austrian Pratincole . . . . .	ii 153
" <i>melanoëtos</i> . . . . .	i 11	Avocet . . . . .	iii 29
" <i>noëvia</i> . . . . .	i 11	Avocet, Common . . . . .	iii 29
Arctic Gull . . . . .	iii 164	" Scooping . . . . .	iii 29



	Vol.	Page		Vol.	Page
Awl-bird . . . . .	i	120	Bittern, Swabian . . . . .	iii	9
<b>B</b>			Bitter Bump . . . . .	iii	11
Baillon's Crake . . . . .	iii	54	Black and White Dabchick . . . . .	iii	110
Baker, Jack . . . . .	i	62	"    "    Wagtail . . . . .	i	142
<i>Balbusardus hatiaëtus</i> . . . . .	i	13	Blackbird . . . . .	ii	53
Bald Coot . . . . .	iii	58	Blackbird, Marsh . . . . .	ii	38
Bank Martin . . . . .	i	141	"    Moor . . . . .	ii	56
Barbary Partridge . . . . .	ii	146	"    Mountain . . . . .	ii	56
Barnacle Goose . . . . .	iii	68	"    Red-winged . . . . .	ii	38
Barn Owl . . . . .	i	54	"    Swamp . . . . .	ii	38
Barred Warbler . . . . .	ii	96	Black Bonnet . . . . .	i	177
Barred Woodpecker . . . . .	i	123	Blackcap . . . . .	ii	97
Bar-tailed Godwit . . . . .	iii	32	Blackcap . . . . .	i	66
Bartram's Sandpiper . . . . .	iii	42	"    . . . . .	i	77
Bartram's Tattler . . . . .	iii	42	"    . . . . .	ii	74
Bay Ibis . . . . .	iii	17	Blackcap Warbler . . . . .	ii	97
Beam Bird . . . . .	i	87	Black-chin Grebe . . . . .	iii	112
Bean Goose . . . . .	iii	65	Black Cock . . . . .	ii	135
Bearded Pinnock . . . . .	i	82	"    Game . . . . .	ii	135
"    Tit . . . . .	i	82	Black Grouse . . . . .	ii	135
Bearded Titmouse . . . . .	i	82	Black Guillemot . . . . .	iii	121
Bee Bird . . . . .	i	87	Black-headed Bunting . . . . .	i	170
Beech-finch . . . . .	ii	1	Black-headed Bunting . . . . .	i	177
Bee-eater . . . . .	i	93	Black-headed Gull . . . . .	iii	148
Bee-eater, Common . . . . .	i	93	Black-headed Snipe . . . . .	iii	22
"    Yellow-throated . . . . .	i	93	Black Kite . . . . .	i	25
Belted Kingfisher . . . . .	i	92	Black Martin . . . . .	i	133
Bent, Joe . . . . .	i	77	"    Neb . . . . .	i	99
Bernacle Goose . . . . .	iii	68	"    Redstart . . . . .	ii	72
Bernicle, Red-breasted . . . . .	iii	69	"    Red-tail . . . . .	ii	72
Bewick's Swan . . . . .	iii	73	"    Sandpiper . . . . .	iii	49
Bilcock . . . . .	iii	55	"    Scoter . . . . .	iii	93
Billy-biter . . . . .	i	72	Blackstart . . . . .	ii	72
Bill, Broad . . . . .	iii	78	Black Stork . . . . .	iii	15
Bird, Beam . . . . .	i	87	Black-tailed Godwit . . . . .	iii	31
"    Bee . . . . .	i	87	Black Tern . . . . .	iii	142
"    Cob-web . . . . .	i	87	Black-throated Diver . . . . .	iii	115
"    Cricket . . . . .	ii	82	Black-throated Grossbeak . . . . .	ii	14
"    Greater Butcher . . . . .	i	59	Black-throated Wheatear . . . . .	ii	78
"    Least . . . . .	i	82	Black-throated Yellow		
"    Lesser . . . . .	i	62	Hammer . . . . .	i	182
"    Ling . . . . .	i	154	Black-toed Gull . . . . .	iii	164
"    Post . . . . .	i	87	"    -winged Longshank . . . . .	iii	30
"    Sedge . . . . .	ii	84	"    "    Stilt . . . . .	iii	30
Bittern . . . . .	iii	11	Black Woodpecker . . . . .	i	119
"    American . . . . .	iii	13	Blue-back . . . . .	ii	45
Bittern, Common . . . . .	iii	11	"    -bonnet . . . . .	i	72
Bittern, Little . . . . .	iii	12	Bluebreast . . . . .	ii	67
Bittern, Mokoho . . . . .	iii	13	Blue-cap . . . . .	i	72
"    Rufous . . . . .	iii	12	"    -headed Wagtail . . . . .	i	148
"    Senegal . . . . .	iii	9	"    Mope . . . . .	i	72
			"    -tail . . . . .	ii	45
			"    -throated Redstart . . . . .	ii	67
			"    "    Robin . . . . .	ii	67



	Vol.	Page		Vol.	Page
Blue-throated Warbler . . .	ii	67	Bunting . . . . .	i	175
“ Tit . . . . .	i	72	“ Black-headed . . .	i	170
Blue Titmouse . . . . .	i	72	Bunting, Black-headed .	i	177
“ -winged Teal . . . .	iii	85	Bunting, Cirl . . . . .	i	182
Bog Bumper . . . . .	iii	11	Bunting, Common . . .	i	175
Bohemian Chatterer . . .	i	113	“ Corn . . . . .	i	175
“ Waxwing . . . . .	i	113	“ Crow . . . . .	i	101
<i>Bombycilla Bohemica</i> . .	i	113	Bunting, Lapland . . .	i	173
“ <i>garrula</i> . . . . .	i	113	Bunting, Lapland Lark .	i	173
<i>Bombycivora garrula</i> . .	i	113	“ Lark . . . . .	i	175
Bonnet, Black . . . . .	i	177	Bunting, Little . . . . .	i	179
Bonxie . . . . .	iii	161	Bunting, Oortolan . . .	i	183
Boonk . . . . .	iii	12	“ Passerine . . . . .	i	177
<i>Botauris freti Hudsonis</i> .	iii	13	Bunting, Reed . . . . .	i	177
“ <i>pusillus</i> . . . . .	iii	12	“ Rustic . . . . .	i	179
“ <i>stellaris</i> . . . . .	iii	11	“ Snow . . . . .	i	171
Bottle Tit . . . . .	i	79	Bunting, Tawny . . . .	i	171
“ Tom . . . . .	i	79	“ Yellow . . . . .	i	180
<i>Brachylopus murarius</i> . .	i	133	Buonaparte's Gull . . .	iii	146
<i>Brachylophus viridis</i> . .	i	120	Buonapartian Gull . . .	iii	146
Bramble Finch . . . . .	ii	4	Burrow Duck . . . . .	iii	76
Brambling . . . . .	i	171	Bustard, Great . . . . .	ii	151
“ . . . . .	ii	4	Bustard, Lesser . . . .	ii	152
“ Greater . . . . .	i	171	Bustard, Little . . . . .	ii	152
Bran-tail . . . . .	ii	69	“ Macqueen's . . . . .	ii	154
“ Fiery . . . . .	ii	69	Bustard, Thick-kneed .	ii	155
Brent Goose . . . . .	iii	68	<i>Buteo cineraceus</i> . . . .	i	47
Bridled Guillemot . . . .	iii	120	“ <i>lagopus</i> . . . . .	i	18
Brighton Pipit . . . . .	i	153	“ <i>nissus</i> . . . . .	i	41
Broad Bill . . . . .	iii	78	“ <i>palumbarius</i> . . . . .	i	39
Broad-billed Sandpiper . .	iii	44	“ <i>vulgaris</i> . . . . .	i	16
Brook-runner . . . . .	iii	55	<i>Butor minutus</i> . . . . .	iii	12
Brown Gull . . . . .	iii	161	Buzzard . . . . .	i	16
Brown-headed Gull . . . .	iii	147	“ Honey . . . . .	i	20
“ Linnet . . . . .	ii	20	Buzzard, Moor . . . . .	i	44
“ Longbeak . . . . .	iii	39	Buzzard, Rough-legged .	i	18
“ Owl . . . . .	i	53			
“ Sandpiper . . . . .	iii	45			
Brown Snipe . . . . .	iii	39			
Brunnich's Guillemot . . .	iii	119			
<i>Bubo maximus</i> . . . . .	i	50			
“ <i>scops</i> . . . . .	i	51			
Buffon's Skua . . . . .	iii	164			
Buff-backed Heron . . . . .	iii	8			
Buff-breasted Sandpiper . .	iii	43			
<i>Budytes flava</i> . . . . .	i	148			
“ <i>Rayi</i> . . . . .	i	149			
Bullfinch . . . . .	ii	26			
Bullfinch, Common . . . .	ii	26			
“ Pine . . . . .	ii	29			
Bullfinch, Scarlet . . . . .	ii	28			
Bulwer's Petrel . . . . .	iii	67			
Bump, Bitter . . . . .	iii	11			
Bumper, Bog . . . . .	iii	11			

## C

<i>Calamoherpe phragmitis</i> .	ii	84
<i>Calamophilus biarmicus</i> .	i	82
<i>Calidris arenaria</i> . . . .	ii	171
“ <i>Canutus</i> . . . . .	iii	41
Calloo . . . . .	iii	100
Canada Goose . . . . .	iii	71
Canada Owl . . . . .	i	58
<i>Cancrofagus luteus</i> . . . .	iii	9
“ <i>minutus</i> . . . . .	iii	11
“ <i>rufus</i> . . . . .	iii	9
Capercaillie . . . . .	ii	133
<i>Caprimulgus Europæus</i> . .	i	131
“ <i>punctatus</i> . . . . .	i	131
“ <i>ruficollis</i> . . . . .	i	130



	Vol.	Page		Vol.	Page
<i>Carlo cormoranus</i> . . .	iii	129	Chopper, Cherry . . .	i	87
“ <i>cristatus</i> . . .	iii	130	Chough . . .	i	95
<i>Carduelis elegans</i> . . .	ii	16	Chough, Cornish . . .	i	95
“ <i>spinus</i> . . .	ii	18	<i>Chroicocephalus capistratus</i> . . .	iii	147
Carolina Cuckoo . . .	i	126	Church Owl . . .	i	54
Carrion Crow . . .	i	99	Churn Owl . . .	i	131
<i>Caryocatactes nucifraga</i> . . .	i	110	<i>Ciconia alba</i> . . .	iii	14
Caspian Tern . . .	iii	132	“ <i>nigra</i> . . .	iii	15
Castaneous Duck . . .	iii	97	<i>Cinclus aquaticus</i> . . .	ii	40
“ Heron . . .	iii	9	Cinereous Godwit . . .	iii	28
<i>Cataracta cephus</i> . . .	iii	164	“ Shrike . . .	i	59
“ <i>parasiticus</i> . . .	iii	164	<i>Circus cyaneus</i> . . .	i	46
“ <i>pomerinus</i> . . .	iii	162	“ <i>Montagui</i> . . .	i	47
“ <i>skua</i> . . .	iii	161	“ <i>rufus</i> . . .	i	44
“ <i>vulgaris</i> . . .	iii	161	Girl Bunting . . .	i	182
<i>Catharacta skua</i> . . .	iii	161	<i>Clangula chrysophthalmus</i> . . .	iii	102
<i>Cathartes percnopterus</i> . . .	i	3	“ <i>glacialis</i> . . .	iii	100
<i>Cenchnis cenchris</i> . . .	i	38	“ <i>histrionica</i> . . .	iii	101
<i>Cephus grylle</i> . . .	iii	121	“ <i>vulgaris</i> . . .	iii	102
<i>Certhia familiaris</i> . . .	i	117	Cliff Daw . . .	i	95
Chack-bird . . .	ii	80	Climber, Tree . . .	i	117
Chaff, Chiff . . .	ii	112	Clod-hopper . . .	ii	80
Chaffinch . . .	ii	1	Coal-and-candle-light . . .	iii	100
Chanchider . . .	i	87	Cobb . . .	iii	159
<i>Charadrius Africanus</i> . . .	ii	157	Cobbler's-awl Duck . . .	iii	29
“ <i>albifrons</i> . . .	ii	163	Cob-web Bird . . .	i	87
“ <i>Alexandrinus</i> . . .	ii	163	<i>Coccothraustes Chloris</i> . . .	ii	11
“ <i>calidris</i> . . .	ii	171	“ <i>vulgaris</i> . . .	ii	14
“ <i>Cantianus</i> . . .	ii	163	<i>Coccyzus Americanus</i> . . .	i	126
“ <i>crepitans</i> . . .	ii	155	Cock, Black . . .	ii	135
“ <i>hiaticula</i> . . .	ii	160	Coldfinch . . .	i	84
“ <i>himantopus</i> . . .	iii	30	Colemouse . . .	i	69
“ <i>littoralis</i> . . .	ii	163	Cole Tit . . .	i	69
“ <i>minor</i> . . .	ii	162	Cole Titmouse . . .	i	69
“ <i>morinellus</i> . . .	ii	159	Collarded Pratincole . . .	ii	153
“ <i>œdicnemus</i> . . .	ii	155	“ Stare . . .	ii	59
“ <i>pluvialis</i> . . .	ii	157	Collier, Tom . . .	i	66
“ <i>rubidus</i> . . .	ii	171	<i>Columba ænas</i> . . .	ii	126
“ <i>Sibiricus</i> . . .	ii	159	“ <i>livia</i> . . .	ii	128
“ <i>Tartaricus</i> . . .	ii	159	“ <i>migratoria</i> . . .	ii	130
Chatterer, Bohemian . . .	i	113	“ <i>palumbus</i> . . .	ii	124
“ European . . .	i	113	“ <i>Turtur</i> . . .	ii	129
“ Waxen . . .	i	113	<i>Colymbus arcticus</i> . . .	iii	115
Chauk Daw . . .	i	95	“ <i>auritus</i> . . .	iii	111
<i>Chauliodus strepera</i> . . .	iii	79	“ <i>borealis</i> . . .	iii	116
Cheeter . . .	i	62	“ <i>Caspicus</i> . . .	iii	110
<i>Chenalopec Egyptiaca</i> . . .	iii	70	“ <i>cornutus</i> . . .	iii	108
<i>Chen Hyperboreus</i> . . .	iii	69	“ . . .	iii	110
Cherry Chopper . . .	i	87	“ <i>cristatus</i> . . .	iii	108
“ Sucker . . .	i	87	“ <i>fluvialis</i> . . .	iii	112
Chicken, Mother Carey's . . .	iii	170	“ <i>gracialis</i> . . .	iii	114
Chiff Chaff . . .	ii	112	“ <i>Hebridicus</i> . . .	iii	112
Chimney Swallow . . .	i	136	“ <i>Immer</i> . . .	iii	114
Chink . . .	i	177	“ <i>minor</i> . . .	iii	112



	Vol.	Page		Vol.	Page
<i>Colymbus obscurus</i> . . .	iii	110	Common Swallow . . .	i	136
" <i>septentrionalis</i> . . .	iii	116	" Swift . . .	i	133
" <i>stellaris</i> . . .	iii	116	" Teal . . .	iii	86
" <i>striatus</i> . . .	iii	116	" Thick-knee . . .	ii	155
Common Avocet . . .	iii	29	" Thrush . . .	ii	49
" Bee-eater . . .	i	93	" Turnstone . . .	ii	167
" Bernicle Goose . . .	iii	68	" Whitethroat . . .	ii	103
" Bittern . . .	iii	11	" Wild Duck . . .	iii	81
" Bullfinch . . .	ii	26	" Wren . . .	ii	116
" Bunting . . .	i	176	Coot . . .	iii	58
" Coot . . .	iii	58	Coot, Bald . . .	iii	58
" Cormorant . . .	iii	129	" Common . . .	iii	58
" Coulter-neb . . .	iii	124	Coot-foot, Red-necked . . .	iii	62
" Crane . . .	iii	1	<i>Coracias garrula</i> . . .	i	89
" Creeper . . .	i	117	Corbie . . .	i	97
" Crossbill . . .	ii	30	Corbie-crow . . .	i	97
" Cuckoo . . .	i	128	" " Great . . .	i	97
" Curlew . . .	iii	19	Cormorant . . .	iii	129
" Dipper . . .	ii	40	Cormorant, Common . . .	iii	129
" Gadwall . . .	iii	79	" Crested . . .	iii	130
" Gallinule . . .	iii	56	" Green . . .	iii	130
" Gannet . . .	iii	131	Cormorant, Green . . .	iii	130
" Godwit . . .	iii	31	Corn Bunting . . .	i	176
" " . . .	iii	32	Corn-crake . . .	iii	50
" Golden-eye . . .	iii	102	Cornish Chough . . .	i	95
" Grosbeak . . .	ii	14	" Daw . . .	i	95
" Guillemot . . .	iii	117	<i>Cornix frugilega</i> . . .	i	103
Common Gull . . .	iii	156	" <i>nigra</i> . . .	i	103
Common Hawfinch . . .	ii	29	Corn Thief . . .	ii	38
" Heron . . .	iii	3	Cornwall Kae . . .	i	95
" Hoopoe . . .	i	94	Corvorant . . .	iii	129
" Kingfisher . . .	i	90	" . . .	iii	130
" Kingfisher . . .	i	90	<i>Corvus caryocatactes</i> . . .	i	110
" Lapwing . . .	ii	165	" <i>cinerea</i> . . .	i	101
" Linnet . . .	ii	20	" <i>corax</i> . . .	i	97
" Magpie . . .	i	108	" <i>cornix</i> . . .	i	101
" Partridge . . .	ii	142	" <i>corone</i> . . .	i	99
" Pheasant . . .	ii	131	" <i>docilis</i> . . .	i	95
" Pintail . . .	iii	80	" <i>frugilegus</i> . . .	i	103
" Pochard . . .	iii	96	" <i>graculus</i> . . .	i	95
" Quail . . .	ii	149	" <i>glandarius</i> . . .	i	111
" Redshank . . .	iii	23	" <i>monedula</i> . . .	i	105
" Rotche . . .	iii	123	" <i>pica</i> . . .	i	108
Common Sandpiper . . .	iii	26	<i>Corythus enculeator</i> . . .	ii	29
Common Scoter . . .	iii	93	Coulter-neb . . .	iii	124
" Scraber . . .	iii	121	" Common . . .	iii	124
" Shieldrake . . .	iii	76	Courland Snipe . . .	iii	22
" Shoveler . . .	iii	78	Courser . . .	ii	154
" Skua . . .	iii	161	Cowbird . . .	i	126
" Snipe . . .	iii	36	Crake, Baillon's . . .	iii	54
" Sparrow . . .	ii	9	" Little . . .	iii	53
" Stare . . .	ii	35	Crake, Olivaceous . . .	iii	53
" StormPetrel . . .	iii	170	Crake, Spotted . . .	iii	51
" Starling . . .	ii	35	Cran . . .	i	133



	Vol.	Page		Vol.	Page
Crane . . . . .	iii	1	Cuckoo's Maid . . . . .	i	116
Crane, Common . . . . .	iii	1	Cuckoo's Mate . . . . .	i	119
Crane, Numdian . . . . .	iii	2	Cuckoo's Messenger . . . . .	i	116
Crank-bird . . . . .	i	123	Cuckoo, Virginian . . . . .	i	126
Cravat Goose . . . . .	iii	71	<b>Cuckoo, Yellow-billed</b> . . . . .	i	126
<b>Creeper</b> . . . . .	i	117	<i>Cuculus Americanus</i> . . . . .	i	126
Creeper, Common . . . . .	i	117	“ <i>canorus</i> . . . . .	i	128
“ Familiar . . . . .	i	117	“ “ <i>rufus</i> . . . . .	i	128
“ Grey . . . . .	i	154	“ <i>Carolinensis</i> . . . . .	i	126
“ Moss . . . . .	i	154	“ <i>cinerosus</i> . . . . .	i	126
“ Tree . . . . .	i	117	“ <i>glandarius</i> . . . . .	i	130
<b>Creeper, Wall</b> . . . . .	i	118	“ <i>hepaticus</i> . . . . .	i	128
Crested Cormorant . . . . .	iii	130	<b>Curlew</b> . . . . .	iii	19
“ Grebe . . . . .	iii	108	Curlew, Common . . . . .	iii	19
“ Heron . . . . .	iii	4	“ Esquimaux . . . . .	iii	20
“ Lapwing . . . . .	ii	165	“ Pigmy . . . . .	iii	40
“ Purple Heron . . . . .	iii	4	<b>Curlew Sandpiper</b> . . . . .	iii	40
<b>Crested Lark</b> . . . . .	i	168	Curlew, Stone . . . . .	ii	155
Crested Shag . . . . .	iii	130	“ Whimbrel . . . . .	iii	20
“ Tit . . . . .	i	70	<i>Curruca arundinacea</i> . . . . .	ii	86
<b>Crested Titmouse</b> . . . . .	i	70	“ <i>atricapilla</i> . . . . .	ii	97
<i>Crex Baillonii</i> . . . . .	iii	54	“ <i>cinerea</i> . . . . .	ii	103
“ <i>porzana</i> . . . . .	iii	51	“ <i>garrula</i> . . . . .	ii	105
“ <i>pratensis</i> . . . . .	iii	50	“ <i>hortensis</i> . . . . .	ii	101
“ <i>pusilla</i> . . . . .	iii	53	“ <i>locustella</i> . . . . .	ii	82
Cricket Bird . . . . .	ii	82	“ <i>lusciniæ</i> . . . . .	ii	90
Crimson-headed Finch . . . . .	ii	28	“ <i>Orphea</i> . . . . .	ii	99
Crooked Bill . . . . .	iii	29	“ <i>sepiaria</i> . . . . .	ii	60
<b>Crossbill</b> . . . . .	ii	30	“ <i>sibilatrix</i> . . . . .	ii	107
Crossbill, Common . . . . .	ii	30	“ <i>Suecica</i> . . . . .	ii	67
“ European . . . . .	ii	30	“ <i>sylvia</i> . . . . .	ii	103
<b>Crossbill, Parrot</b> . . . . .	ii	32	“ <i>sylviella</i> . . . . .	ii	105
“ White-winged . . . . .	ii	33	<i>Cursorius Europæus</i> . . . . .	ii	154
<b>Crow</b> . . . . .	i	99	Curwillet . . . . .	ii	171
Crow, Bunting . . . . .	i	101	Cushat . . . . .	ii	124
“ Carrion . . . . .	i	99	<i>Cygnus Bewickii</i> . . . . .	iii	73
“ Dun . . . . .	i	101	“ <i>Canadensis</i> . . . . .	iii	71
“ Flesh . . . . .	i	99	“ <i>ferus</i> . . . . .	iii	72
“ Gesner's Wood . . . . .	i	95	“ <i>mansuetus</i> . . . . .	iii	74
“ Gor . . . . .	i	99	“ <i>olor</i> . . . . .	iii	74
“ Gore . . . . .	i	99	<i>Cypselus alpinus</i> . . . . .	i	135
“ Grey . . . . .	i	101	“ <i>apus</i> . . . . .	i	133
“ Grey-backed . . . . .	i	101	“ <i>murarius</i> . . . . .	i	133
“ Hermit . . . . .	i	95			
<b>Crow, Hooded</b> . . . . .	i	101			
<b>Crow, Market-jew</b> . . . . .	i	95			
“ Red-legged . . . . .	i	95			
“ Royston . . . . .	i	101			
“ Water . . . . .	ii	40			
<b>Cuckoo</b> . . . . .	i	128			
Cuckoo, American . . . . .	i	126			
“ Carolina . . . . .	i	126			
“ Common . . . . .	i	128			
<b>Cuckoo, Great Spotted</b> . . . . .	i	130			



	Vol.	Page		Vol.	Page
Dalmatian Regulus . . .	ii	123	Duck, Red . . .	iii	97
Dartford Warbler . . .	ii	114	“ Red-crested . . .	iii	95
Daw . . .	i	105	Duck, Red-crested Whistling . . .	iii	95
“ Chalk . . .	i	95	Duck, Scaup . . .	iii	98
“ Cliff . . .	i	95	“ Sharp-tailed . . .	iii	100
“ Cornish . . .	i	44	“ Shoveler . . .	iii	78
Desert Wheatear . . .	ii	79	“ St. Cuthbert's . . .	iii	89
Didapper . . .	iii	112	Duck, Tufted . . .	iii	99
Dipper . . .	ii	40	Duck, Western . . .	iii	88
Dipper, Common . . .	ii	40	“ White-eyed . . .	iii	97
“ European . . .	ii	40	Duck, Wild . . .	iii	81
Dish-washer . . .	i	142	Dull-willy . . .	ii	160
Diver, Black-throated . . .	iii	115	Dun Crow . . .	i	101
Diver, Dun . . .	iii	106	“ Diver . . .	iii	106
“ First Speckled . . .	iii	116	Dunkadoo . . .	iii	13
“ Greatest Speckled . . .	iii	114	Dunlin . . .	iii	48
Diver, Great Northern . . .	iii	114	Dunlin, Minute . . .	iii	45
Diver, Imber . . .	iii	116	“ Sandpiper . . .	iii	48
“ Lough . . .	iii	103	“ Temminck's . . .	iii	46
“ Northern . . .	iii	114	Duncock . . .	ii	60
Diver, Red-throated . . .	iii	116	Dusky Grebe . . .	iii	110
Diver, Speckled . . .	iii	116	Dusky Lark . . .	i	160
“ Striped . . .	iii	116	“ Sandpiper . . .	iii	22
Dor-hawk . . .	i	131	“ Snipe . . .	iii	22
Dotterel . . .	ii	159	Dwarf Heron . . .	iii	9
“ Kentish . . .	ii	163	“ Rail . . .	iii	53
“ Little Ringed . . .	ii	162			
Dotterel Plover . . .	ii	159			
Dotterel, Ringed . . .	ii	160			
Dottrel . . .	ii	159			
“ Little Ringed . . .	ii	162			
Double Snipe . . .	iii	35			
Doucker, Great . . .	iii	114			
“ Northern . . .	iii	115			
“ Small . . .	iii	112			
Dove, Greenland . . .	iii	121			
“ Ring . . .	ii	124			
Dove, Rock . . .	ii	128			
“ Stock . . .	ii	126			
“ Turtle . . .	ii	129			
Duck, Burrow . . .	iii	76			
“ Castaneous . . .	iii	97			
“ Common Wild . . .	iii	81			
Duck, Eider . . .	iii	98			
“ Ferruginous . . .	iii	97			
Duck, Gadwall . . .	iii	79			
“ Gargany . . .	iii	83			
“ Golden-eye . . .	iii	102			
“ Harlequin . . .	iii	101			
“ Hawk . . .	i	44			
Duck, King . . .	iii	91			
“ Long-tailed . . .	iii	100			
Duck, Pintail . . .	iii	80			
“ Pochard . . .	iii	96			

## E

<i>Eagle fulvus</i> . . .	i	8
Eagle, Golden . . .	i	8
Eagle Owl . . .	i	50
“ Spotted . . .	i	11
Eared Dabchick . . .	iii	111
Eared Grebe . . .	iii	111
Ecle . . .	i	120
<i>Ectopistes migratorius</i> . . .	ii	130
Egret, Great . . .	iii	6
“ “ White . . .	iii	6
“ Heron . . .	iii	7
Egret, Little . . .	iii	7
Egret, Rufous-backed . . .	iii	8
Egyptian Goose . . .	iii	70
“ Vulture . . .	i	3
Eider Duck . . .	iii	89
Eider, King . . .	iii	91
Eider, Stellar's . . .	iii	88
Eider, Western . . .	iii	88
<i>Elanus furcatus</i> . . .	i	24
Elk . . .	iii	72
<i>Emberiza arundinacea</i> . . .	i	177
“ <i>chlorocephala</i> . . .	i	183



	Vol.	Page		Vol.	Page
<i>Emberiza citrlus</i> . . .	i	182	<i>Falco naevius</i> . . .	i	11
“ <i>citrinella</i> . . .	i	180	“ <i>nigrans</i> . . .	i	25
“ <i>elcathorax</i> . . .	i	182	“ <i>nisus</i> . . .	i	41
“ <i>flava</i> . . .	i	180	“ <i>palumbarius</i> . . .	i	39
“ <i>glacialis</i> . . .	i	171	“ <i>peregrinus</i> . . .	i	28
“ <i>hortulana</i> . . .	i	183	“ <i>rufipes</i> . . .	i	32
“ <i>lapponica</i> . . .	i	173	“ <i>subbuteo</i> . . .	i	30
“ <i>melanocephala</i> . . .	i	170	“ <i>Tinnuncularis</i> . . .	i	38
“ <i>miliaria</i> . . .	i	175	“ <i>Tinnunculoides</i> . . .	i	38
“ <i>montana</i> . . .	i	171	“ <i>Tinnunculus</i> . . .	i	35
“ <i>mustelina</i> . . .	i	171	“ <i>vespertinus</i> . . .	i	32
“ <i>nivalis</i> . . .	i	171	Fallow-chat . . .	ii	80
“ <i>passerina</i> . . .	i	177	Familiar Creeper . . .	i	117
“ <i>pusilla</i> . . .	i	179	Fauvette, Reed . . .	ii	84
“ <i>rustica</i> . . .	i	179	“ Winter . . .	ii	60
“ <i>schœnielus</i> . . .	i	177	<i>Fedoa melanura</i> . . .	iii	31
“ <i>schœniculus</i> . . .	i	177	“ <i>Meyeri</i> . . .	iii	32
“ <i>Tunstalli</i> . . .	i	183	“ <i>rufa</i> . . .	iii	32
Emmet-hunter . . .	i	116	Feldfare . . .	ii	45
Epicurean Warbler . . .	i	84	Felt . . .	ii	45
Equestrian Sandpiper . . .	iii	33	Feltfare . . .	ii	45
Erne . . .	i	6	Fern-owl . . .	i	131
<i>Erythaca rubecula</i> . . .	ii	62	Ferruginous Duck . . .	iii	97
<i>Erythrophrys Americanus</i> . . .	i	126	<i>Ficedula Suecica</i> . . .	ii	67
Esquimaux Curlew . . .	iii	20	“ <i>Tiathy's</i> . . .	ii	72
European Chatterer . . .	i	113	Fieldfare . . .	ii	45
“ Crossbill . . .	ii	30	Field Lark . . .	i	158
“ Dipper . . .	ii	40	“ “ . . .	i	160
“ Goat-sucker . . .	i	131	“ “ . . .	i	167
			“ Titling . . .	i	158
			Fiery Bran-tail . . .	ii	69
			Fighting Ruff . . .	iii	33
			Finch, Bramble . . .	ii	4
			“ Crimson-headed . . .	ii	28
			“ Great Pied Mountain . . .	i	171
			“ Lapland . . .	i	173
			“ Lesser Mountain . . .	i	171
			“ Lulean . . .	ii	4
			Finch, Mountain . . .	ii	4
			“ Serin . . .	ii	13
			Firecrest . . .	ii	122
			Fire-crested Kinglet . . .	ii	122
			“ “ Wren . . .	ii	122
			“ -crowned Kinglet . . .	ii	122
			Fire-tail . . .	ii	69
			First Speckled Diver . . .	iii	116
			Flake, Snow . . .	i	171
			Flamingo . . .	iii	18
			Fleck . . .	i	171
			Flesh Crow . . .	i	99
			Flusher . . .	i	62
			Flycatcher, Pied . . .	i	84
			“ Red-breasted . . .	i	86

## F

<i>Falco æruginosus</i> . . .	i	44
“ <i>æsalon</i> . . .	i	33
“ <i>albicilla</i> . . .	i	6
“ <i>apivorus</i> . . .	i	20
“ <i>arundinaceus</i> . . .	i	44
“ <i>buteo</i> . . .	i	16
“ <i>cenchris</i> . . .	i	38
“ <i>chrysoëtos</i> . . .	i	8
“ <i>cineraceus</i> . . .	i	47
“ <i>cinerareus</i> . . .	i	47
“ <i>cinerarius</i> . . .	i	47
“ <i>communis</i> . . .	i	28
“ <i>fureatus</i> . . .	i	24
“ <i>gracilis</i> . . .	i	38
“ <i>Gyrfalco</i> . . .	i	26
“ <i>haliaëtus</i> . . .	i	13
“ <i>hyemalis</i> . . .	i	47
“ <i>Islandicus</i> . . .	i	26
“ <i>lagopus</i> . . .	i	18
“ <i>maculatus</i> . . .	i	11
“ <i>milvus</i> . . .	i	22



	Vol.	Page
<b>Flycatcher, Spotted</b> . . .	i	87
Foolish Guillemot . . .	iii	117
Fork-tailed Kite . . .	i	22
<b>Fork-tailed Petrel</b> . . .	iii	169
“ “ Storm Petrel . . .	iii	169
Fowl, Snow . . .	i	171
Freckled Heron . . .	iii	13
<i>Fregilus graculus</i> . . .	i	95
French Partridge . . .	ii	145
“ Pie . . .	i	122
“ Pigeon . . .	ii	165
“ Woodpecker . . .	i	122
“ Yellow-hammer . . .	i	182
<i>Fringilla borealis</i> . . .	ii	24
“ <i>cælebs</i> . . .	ii	1
“ <i>cannabina</i> . . .	ii	20
“ <i>carduelis</i> . . .	ii	16
“ <i>Chloris</i> . . .	ii	11
“ <i>coccothraustes</i> . . .	ii	14
“ <i>domestica</i> . . .	ii	9
“ <i>flavirostris</i> . . .	ii	25
“ <i>linaria</i> . . .	ii	22
“ <i>Linota</i> . . .	ii	20
“ <i>Lulensis</i> . . .	ii	4
“ <i>montana</i> . . .	ii	7
“ <i>montifringilla</i> . . .	ii	4
“ <i>rufescens</i> . . .	ii	22
“ <i>serinus</i> . . .	ii	13
“ <i>spinus</i> . . .	ii	18
<i>Fulica alerrima</i> . . .	iii	58
“ <i>atra</i> . . .	iii	58
“ <i>chloropus</i> . . .	iii	56
<i>Fuligula cristata</i> . . .	iii	99
“ <i>dispar</i> . . .	iii	88
“ <i>ferina</i> . . .	iii	96
“ <i>marila</i> . . .	iii	98
“ <i>nyroca</i> . . .	iii	97
“ <i>rufina</i> . . .	iii	95
<b>Fulmar</b> . . .	iii	165
Fulmar, Northern . . .	iii	165
“ Petrel . . .	iii	165
“ “ . . .	iii	165
Furzechat . . .	ii	76
Furze Wren . . .	ii	114

## G

<b>Gadwall</b> . . .	iii	79
Gadwall Duck . . .	iii	79
“ Common . . .	iii	79
Gairfowl . . .	iii	128
<i>Galgulus argentoratensis</i> . . .	i	89
“ <i>garrula</i> . . .	i	89

	Vol.	Page
<i>Gallinago major</i> . . .	iii	53
“ <i>media</i> . . .	iii	36
“ <i>minima</i> . . .	iii	38
“ <i>minor</i> . . .	iii	38
<i>Gallinula Baillonii</i> . . .	iii	54
“ <i>chloropus</i> . . .	iii	56
“ <i>crex</i> . . .	iii	50
“ <i>minuta</i> . . .	iii	53
“ <i>porzana</i> . . .	iii	51
“ <i>pusilla</i> . . .	iii	53
Gallinule, Common . . .	iii	56
“ Spotted . . .	iii	51
Gambet Sandpiper . . .	iii	23
Game, Black . . .	ii	135
<b>Gannet</b> . . .	iii	131
Gannet, Common . . .	iii	131
“ Solan . . .	iii	131
Gardenian Heron . . .	iii	10
<b>Garden Warbler</b> . . .	ii	101
<b>Gargeny</b> . . .	iii	83
Garganey Duck . . .	iii	83
“ Teal . . .	iii	83
Garrot . . .	iii	102
<i>Garrulus glandarius</i> . . .	i	111
Garrulus Roller . . .	i	89
Gaunt . . .	iii	108
German Parrot . . .	i	89
Gesner's Wood Crow . . .	i	95
Gibraltar Quail . . .	ii	150
Gilli-howlet . . .	i	54
<i>Glarola Austriaca</i> . . .	ii	153
“ <i>torquata</i> . . .	ii	153
<b>Glaucous Gull</b> . . .	iii	160
Glead . . .	i	22
Glossy Ibis . . .	iii	17
Gnat-snapper . . .	i	93
Goatsucker . . .	i	131
“ European . . .	i	131
“ Nocturnal . . .	i	131
<b>Godwit, Bar-tailed</b> . . .	iii	32
“ <b>Black-tailed</b> . . .	iii	31
Godwit, Cinereous . . .	iii	28
“ Common . . .	iii	31
“ “ . . .	iii	32
“ Grey . . .	iii	32
“ Hudsonian . . .	iii	31
“ Lesser . . .	iii	31
“ Meyer's . . .	iii	32
“ Red . . .	iii	31
“ “ . . .	iii	32
Godwyn . . .	iii	31
“ “ . . .	iii	32
<b>Goldcrest</b> . . .	ii	119
<b>Goldfinch</b> . . .	ii	16



	Vol.	Page		Vol.	Page
Golden-crested Kinglet	ii	119	Great Bustard	ii	151
“ “ Wren	ii	119	Great Corbie Crow	i	97
“ -crowned “	ii	119	Great Crested Grebe	iii	108
Golden Eagle	i	8	Great Doucker	i	114
Golden-eye	iii	102	Great-eared Owl	i	50
Golden-eye, Common	iii	102	Greater Brambling	i	171
“ Duck	iii	102	“ Black-backed Gull	iii	159
Golden Oriole	ii	58	“ Butcher Bird	i	59
Golden Plover	ii	157	“ Loon	iii	108
Goldie	ii	16	“ Pettychaps	ii	101
“ “	i	180	“ Redpole	ii	20
Goldspink	ii	16	“ Speckled Diver	iii	114
Goosander	iii	106	“ Spotted Woodpecker	i	122
Goosander, Green-headed	iii	106	Great Grey Shrike	i	59
“ Red-breasted	iii	105	“ Horned Owl	i	50
“ White-headed	iii	103	Great Northern Diver	iii	114
Goose, Barnacle	iii	68	Great Owl	i	50
Goose, Bean	iii	65	“ Pied Mountain Finch	i	171
“ Bernacle	iii	68	Great Plover	ii	155
“ Brent	iii	68	“ Sedge Warbler	ii	93
“ Canada	iii	71	Great Shrike	i	59
Goose, Common Bernicle	iii	68	Great Snipe	iii	35
“ Cravat	iii	71	“ Spotted Cuckoo	i	130
Goose, Egyptian	iii	70	“ “ Woodpecker	i	122
“ Gray-lag	iii	63	“ Titmouse	i	66
Goose, Grey-lag Wild	iii	63	Great Tit	i	66
“ Laughing	iii	67	“ White Egret	iii	6
Goose, Pink-footed	iii	66	Great White Heron	iii	6
“ Red-breasted	iii	69	Grebe, Black-chin	iii	112
Goose, Siberian	iii	69	“ Crested	iii	108
Goose, Snow	iii	69	Grebe, Dusky	iii	110
Goose, Solan	iii	131	“ Eared	iii	111
“ Soland	iii	131	“ Great Crested	iii	108
“ Wild	iii	65	Grebe, Horned	iii	110
Goose, White-fronted	iii	67	“ Little	iii	112
Goose, White-fronted Wild	iii	67	Grebe, Red-necked	iii	109
Gor-cock	ii	137	Grebe, Sclavonian	iii	110
Gor Crow	i	99	“ Tipper	iii	108
Gore Crow	i	99	Greenfinch	ii	11
Goshawk	i	39	Green Cormorant	iii	130
Gowk	i	128	Green-crested Cormorant	iii	130
Gracula atthis	i	90	Green Grossbeak	ii	11
Grasschat	ii	76	“ Ibis	iii	67
Grasshopper Lark	i	158	Greenland Dove	iii	121
Grasshopper Warbler	ii	82	Green Lapwing	ii	165
Great Auk	iii	128	“ Linnet	ii	11
Great Belted Kingfisher	i	92	“ Plover	ii	157
Great Black-backed Gull	iii	159	“ “	ii	165
“ -headed “	iii	150	Green Sandpiper	iii	24
Great Black-headed Tomtit	i	66	Greenshank	iii	28
“ “ and White Gull	iii	159	Greenwich Sandpiper	iii	33
“ “ “ Woodpecker	i	122	Green Woodpecker	i	120
“ “ Woodpecker	i	119	Green Wren	ii	107



	Vol. Page		Vol. Page
Grey-and-White Wagtail	i 144	Gull, Herring	iii 158
Grey-backed Crow	i 101	Gull, Iceland	iii 155
“ Creeper	i 154	Gull, Ivory	iii 154
“ Crow	i 101	“ Laughing	iii 151
“ Godwit	iii 32	“ Little	iii 146
Grey-headed Wagtail	i 148	“ Lesser Black-backed	iii 157
“ -lag Goose	iii 63	Gull, Lesser White-winged	iii 155
Grey-lag Wild Goose	iii 63	Gull, Masked	iii 147
“ Linnet	ii 20	“ Mediterranean Black-headed	iii 150
Grey Phalarope	iii 61	Gull, Peewit	iii 148
“ Plover	ii 164	“ Red-legged	iii 148
Grey Sandpiper	ii 164	Gull, Sabine's	iii 145
Grey Shrike	i 59	Gull, Silvery	iii 158
Grey Snipe	iii 39	“ Skua	iii 161
Grey Wagtail	i 146	“ White-winged Silvery	iii 155
Griffon Vulture	i 1	“ Yellow-legged	iii 157
Grossbeak	ii 14	<i>Gyps fulvus</i>	i 1
“ Black-throated	ii 14	“ <i>vulgaris</i>	i 1
“ Common	ii 14	<i>Gyr falco candicans</i>	i 26
“ Green	ii 11		
“ Haw	ii 14		
Grossbeak, Pine	ii 29		
Grossbeak, Scarlet	ii 28		
Grouse, Black	ii 135		
“ Red	ii 137		
“ Sand	ii 141		
Grouse, Wood	ii 133		
<i>Grus cinerea</i>	iii 1		
Guillemot	iii 117		
“ Black	iii 121		
Guillemot, Bridled	iii 120		
Guillemot, Brunnich's	iii 119		
Guillemot, Common	iii 117		
“ Foolish	iii 117		
“ Lesser	iii 117		
Guillemot, Ringed	iii 120		
“ Spotted	iii 121		
“ Thick-billed	iii 119		
Gull, Arctic	iii 164		
Gull-billed Tern	iii 140		
“ Black-headed	iii 148		
Gull, Black-toed	iii 164		
“ Brown	iii 161		
“ “ -headed	iii 147		
Gull, Buonaparte's	iii 146		
Gull, Buonapartian	iii 146		
Gull, Common	iii 156		
“ Glaucous	iii 160		
“ Great Black-backed	iii 159		
“ Great Black-headed	iii 150		
Gull, Greater Black and White	iii 159		
“ “ “ -backed	iii 159		
	iii 159		

## H

<i>Hæmatopus Bellonii</i>	ii 172
“ <i>ostralegus</i>	ii 172
Hairy Woodpecker	i 124
Half Snipe	iii 38
<i>Haliaëtus albicilla</i>	i 6
Hareld, Northern	iii 100
Harlequin Duck	iii 101
Harlequin Garrot	iii 101
Harpy, White-headed	i 44
Harrier, Hen	i 46
“ Marsh	i 44
“ Montagu's	i 47
Harrier, Moor	i 44
Harry, King	ii 16
Hawfinch	ii 14
Hawfinch, Common	ii 29
Haw Grossbeak	ii 14
Hawk, Duck	i 44
“ Fishing	i 13
“ Mouse	i 48
“ Owl	i 48
Hawk Owl	i 58
Hawk, Stannel	i 35
Heather-bleater	iii 36
Hebridal Sandpiper	ii 167
Hedge-chanter	ii 60
“ -sparrow	ii 60
“ Warbler	ii 60
Hemipode Andalusian	ii 150



	Vol.	Page		Vol.	Page
<i>Hemipodius lanatus</i> . . . . .	ii	150	<b>Hooded Merganser</b> . . . . .	iii	104
" <i>tachydromus</i> . . . . .	ii	150	Hoody . . . . .	i	101
<b>Hen Harrier</b> . . . . .	i	46	Hoop . . . . .	ii	26
Hermit Crow . . . . .	i	95	<b>Hooper</b> . . . . .	iii	72
Hern . . . . .	iii	3	Hooping Swan . . . . .	iii	72
<b>Heron</b> . . . . .	iii	3	<b>Hoopoe</b> . . . . .	i	94
Heron, African . . . . .	iii	4	Hoopoe, Common . . . . .	i	94
<b>Heron, Buff-backed</b> . . . . .	iii	8	Horned Dabchick . . . . .	iii	110
Heron, Castaneous . . . . .	iii	9	" Grebe . . . . .	iii	110
" Common . . . . .	iii	3	Horsefinch . . . . .	ii	1
" Crested . . . . .	iii	3	Horseman, Red-legged . . . . .	iii	23
" " Purple . . . . .	iii	4	Horsematch . . . . .	i	59
" Dwarf . . . . .	iii	9	House Martin . . . . .	i	139
" Egret . . . . .	iii	7	" Sparrow . . . . .	ii	9
" Freckled . . . . .	iii	13	Howlet . . . . .	i	54
" Gardenian . . . . .	iii	10	Huck-muck . . . . .	i	79
<b>Heron, Great White</b> . . . . .	iii	6	" . . . . .	ii	109
Heron, Jamaica Night . . . . .	iii	10			
" Lesser Ash-coloured . . . . .	iii	10			
" Little . . . . .	iii	12	<b>Ibis</b> . . . . .	iii	17
" " Bittern . . . . .	iii	12	Ibis, Bay . . . . .	iii	17
" " White . . . . .	iii	8	<i>Ibis falcinellus</i> . . . . .	iii	17
<b>Heron, Night</b> . . . . .	iii	10	Ibis, Glossy . . . . .	iii	17
" Purple . . . . .	iii	4	" Green . . . . .	iii	17
Heron, Purple Crested . . . . .	iii	4	<i>Ibis ignea</i> . . . . .	iii	17
" Red-billed . . . . .	iii	8	<b>Iceland Gull</b> . . . . .	iii	155
" Red-legged . . . . .	iii	9	<i>Icterus phænicurus</i> . . . . .	ii	38
Heronsewgh . . . . .	iii	3	Imber Diver . . . . .	iii	114
Heronshaw . . . . .	iii	3	" Lesser . . . . .	iii	115
<b>Heron, Squacco</b> . . . . .	iii	9	Immer . . . . .	iii	114
Heron, Squaiotta . . . . .	iii	9	<b>Isabelline Wheatear</b> . . . . .	ii	78
Hew-hole . . . . .	i	120	<i>Ispida senegalensis</i> . . . . .	i	90
Hickwall . . . . .	i	72	<b>Ivory Gull</b> . . . . .	iii	154
" . . . . .	i	123			
High Hoe . . . . .	i	120			
<i>Himantopus atropterus</i> . . . . .	iii	20			
" <i>melanopterus</i> . . . . .	iii	30			
" <i>Plinii</i> . . . . .	iii	30			
" <i>rufipes</i> . . . . .	iii	30			
<i>Hirundo apus</i> . . . . .	i	133			
" <i>domestica</i> . . . . .	i	136	<b>Jack Baker</b> . . . . .	i	62
" <i>marina</i> . . . . .	ii	153	<b>Jackdaw</b> . . . . .	i	105
" <i>pratincta</i> . . . . .	ii	153	Jackdaw, Red-legged . . . . .	i	95
" <i>purpurea</i> . . . . .	i	138	<b>Jack Snipe</b> . . . . .	iii	38
" <i>riparia</i> . . . . .	i	141	Jadreka Snipe . . . . .	iii	31
" <i>rustica</i> . . . . .	i	136	Jager, Arctic . . . . .	iii	164
" <i>urbica</i> . . . . .	i	139	" Pomerine . . . . .	iii	162
Hissing Owl . . . . .	i	54	Jamaica Night Heron . . . . .	iii	10
<b>Hobby</b> . . . . .	i	30	Jar-Owl . . . . .	i	131
" Orange-legged . . . . .	i	32	<b>Jay</b> . . . . .	i	111
Hoe, High . . . . .	i	120	<b>Jer-Falcon</b> . . . . .	i	26
<b>Honey Buzzard</b> . . . . .	i	20	Jimpo . . . . .	ii	116
<b>Hooded Crow</b> . . . . .	i	101	Joe Bent . . . . .	i	77
			John Whiskey . . . . .	i	62
			Judcock . . . . .	iii	38

## J



## K

	Vol.	Page
Kae . . . . .	i	105
Kae, Cornwall . . . . .	i	95
Kentish Dotterel . . . . .	ii	163
Kentish Plover . . . . .	ii	163
Kestrel . . . . .	i	35
“ Lesser . . . . .	i	38
Killdeer Plover . . . . .	ii	169
Killer, Nine . . . . .	i	62
Killigrew . . . . .	i	95
King Duck . . . . .	iii	91
King Eider . . . . .	iii	91
Kingfisher . . . . .	i	90
“ Belted . . . . .	i	92
Kingfisher, Common . . . . .	i	90
“ Great Belted . . . . .	i	92
King Harry . . . . .	ii	16
Kinglet, Fire-crested . . . . .	ii	122
“ “ -crowned . . . . .	ii	122
“ Golden-crested . . . . .	ii	119
Kingsfisher . . . . .	i	90
“ Common . . . . .	i	90
Kite . . . . .	i	22
“ Black . . . . .	i	25
Kite, Fork-tailed . . . . .	i	22
Kite, Swallow-tailed . . . . .	i	24
Kitty Wren . . . . .	ii	116
Knot . . . . .	iii	41

## L

<i>Lagopus mutus</i> . . . . .	ii	139
“ <i>Scoticus</i> . . . . .	ii	137
“ <i>vulgaris</i> . . . . .	ii	139
Land-rail . . . . .	iii	50
<i>Lanius collurio</i> . . . . .	i	62
“ <i>excubitor</i> . . . . .	i	59
“ <i>minor</i> . . . . .	i	61
“ <i>pomeranus</i> . . . . .	i	64
“ <i>rufus</i> . . . . .	i	64
“ <i>rutilus</i> . . . . .	i	64
Lapland Bunting . . . . .	i	173
Lapland Finch . . . . .	i	173
“ Lark . . . . .	i	173
Lapwing . . . . .	ii	165
“ Common . . . . .	ii	165
“ Crested . . . . .	ii	165
“ Green . . . . .	ii	165
“ Sandpiper . . . . .	ii	165
Large Green Woodpecker . . . . .	i	120
Larger Willow Wren . . . . .	ii	107
Lark, Bunting . . . . .	i	175
Lark, Crested . . . . .	i	168

Vol. Page

Lark, Dusky . . . . .	i	160
“ Field . . . . .	i	160
“ “ . . . . .	i	158
“ Field . . . . .	i	167
“ Grasshopper . . . . .	i	158
“ Lesser Crested . . . . .	i	158
“ “ Field . . . . .	i	158
“ Meadow . . . . .	i	154
“ “ . . . . .	i	158
“ Pipit . . . . .	i	158
“ Rock . . . . .	i	160
“ Sand . . . . .	ii	160
“ Sea . . . . .	i	160
“ “ . . . . .	iii	48
“ Shore . . . . .	i	160
Lark, Shore . . . . .	i	162
Lark, Short-heeled Field . . . . .	i	158
Lark, Short-toed . . . . .	i	164
“ Sky . . . . .	i	167
Lark, Tawny . . . . .	i	152
“ Tree . . . . .	i	158
Lark, White-winged . . . . .	i	162
“ Wood . . . . .	i	165
<i>Larus arcticus</i> . . . . .	iii	155
“ <i>argentatus</i> . . . . .	iii	155
“ “ . . . . .	iii	158
“ <i>atricilla</i> . . . . .	iii	151
“ <i>atricilloides</i> . . . . .	iii	146
“ <i>Buonapartii</i> . . . . .	iii	146
“ <i>candidus</i> . . . . .	iii	154
“ <i>canus</i> . . . . .	iii	156
“ <i>capistratus</i> . . . . .	iii	147
“ <i>cataractes</i> . . . . .	iii	161
“ <i>crepidatus</i> . . . . .	iii	164
“ <i>eburneus</i> . . . . .	iii	154
“ <i>fuscus</i> . . . . .	iii	156
“ “ . . . . .	iii	157
“ <i>glaucoides</i> . . . . .	iii	155
“ <i>glaucus</i> . . . . .	iii	158
“ “ . . . . .	iii	160
“ <i>hybernus</i> . . . . .	iii	156
“ <i>Islandicus</i> . . . . .	iii	155
“ <i>ichthyaetus</i> . . . . .	iii	150
“ <i>leucopterus</i> . . . . .	iii	155
“ <i>marinus</i> . . . . .	iii	158
“ “ . . . . .	iii	159
“ <i>melanocephalus</i> . . . . .	iii	150
“ <i>minutus</i> . . . . .	iii	146
“ <i>naevius</i> . . . . .	iii	159
“ <i>niger</i> . . . . .	iii	159
“ <i>parasiticus</i> . . . . .	iii	163
“ <i>rubundus</i> . . . . .	iii	148
“ <i>rissa</i> . . . . .	iii	152
“ <i>Sabinei</i> . . . . .	iii	145



	Vol.	Page		Vol.	Page
<i>Larus Sabini</i> . . . . .	iii	145	<i>Linaria rubra</i> . . . . .	ii	22
" <i>tridactylus</i> . . . . .	iii	152	Ling Bird . . . . .	i	154
Laughing Goose . . . . .	iii	67	<b>Linnet</b> . . . . .	ii	20
Laughing Gull . . . . .	iii	151	Linnet, Brown . . . . .	ii	20
Lavrock . . . . .	i	167	" Common . . . . .	ii	20
" Sand . . . . .	ii	160	" Green . . . . .	ii	11
" " . . . . .	iii	26	" Grey . . . . .	ii	20
Leach's Petrel . . . . .	iii	169	" Lesser Redpole . . . . .	ii	22
Least Butcher Bird . . . . .	i	82	" " " . . . . .	ii	24
" Snipe . . . . .	iii	45	" Mountain . . . . .	ii	25
" " . . . . .	iii	48	" Red-breasted . . . . .	ii	20
" Spotted Woodpecker . . . . .	i	123	" Rose . . . . .	ii	20
" Willow Wren . . . . .	ii	112	" Whin . . . . .	ii	20
Lesser Ash-coloured Heron . . . . .	iii	10	<i>Linota cannabina</i> . . . . .	ii	20
" Bustard . . . . .	ii	152	" <i>linaria</i> . . . . .	ii	22
" Butcher Bird . . . . .	i	62	Little Auk . . . . .	iii	123
" Crested Lark . . . . .	i	158	<b>Little Bittern</b> . . . . .	iii	12
" Field Lark . . . . .	i	158	Little Bittern Heron . . . . .	iii	12
" Godwit . . . . .	iii	31	" Black and White		
<b>Lesser Grey Shrike</b> . . . . .	i	61	Woodpecker . . . . .	i	123
Lesser Guillemot . . . . .	iii	117	<b>Little Bunting</b> . . . . .	i	179
" Imber . . . . .	iii	115	" <b>Bustard</b> . . . . .	ii	152
<b>Lesser Kestrel</b> . . . . .	i	38	" <b>Crake</b> . . . . .	iii	53
Lesser Mountain Finch . . . . .	i	171	" <b>Egret</b> . . . . .	iii	7
" Pettychaps . . . . .	ii	112	Little French Woodpecker . . . . .	i	123
" Redpole . . . . .	ii	22	" Gallinule . . . . .	iii	53
" " . . . . .	ii	24	" Grebe . . . . .	iii	112
" " Linnet . . . . .	ii	22	<b>Little Gull</b> . . . . .	iii	146
" " " . . . . .	ii	24	Little Heron . . . . .	iii	12
<b>Lesser Spotted Woodpecker</b> . . . . .	i	123	" Horned Owl . . . . .	i	51
" <b>Tern</b> . . . . .	iii	141	" Night Owl . . . . .	i	56
" <b>Whitethroat</b> . . . . .	ii	105	<b>Little Owl</b> . . . . .	i	56
<i>Estris crepidatus</i> . . . . .	iii	164	Little Peep . . . . .	iii	45
" <i>parasiticus</i> . . . . .	iii	163	" Ring Dottrell . . . . .	ii	162
" " . . . . .	iii	164	<b>Little Ringed Dotterel</b> . . . . .	ii	162
" <i>pomerinus</i> . . . . .	iii	162	Little Ringed Plover . . . . .	ii	162
" <i>Richardsonii</i> . . . . .	iii	163	" Sandpiper . . . . .	iii	45
<i>Limosa ægocephala</i> . . . . .	iii	31	" " . . . . .	iii	46
" <i>glottis</i> . . . . .	iii	28	" Stint . . . . .	iii	45
" <i>grisea</i> . . . . .	iii	28	" Tern . . . . .	iii	141
" " <i>major</i> . . . . .	iii	32	" Whaap . . . . .	iii	20
" <i>melanura</i> . . . . .	iii	31	" White Heron . . . . .	iii	8
" <i>rufa</i> . . . . .	iii	32	<i>Lobipes hyperborea</i> . . . . .	iii	62
" " <i>major</i> . . . . .	iii	31	" <i>hyperboreus</i> . . . . .	iii	62
<i>Linaria alnorum</i> . . . . .	ii	22	<i>Locustella avicula</i> . . . . .	ii	82
" <i>betularum</i> . . . . .	ii	22	<b>Long-eared Owl</b> . . . . .	i	49
" <i>borealis</i> . . . . .	ii	24	Long-horned Owl . . . . .	i	49
" <i>canescens</i> . . . . .	ii	24	Long-legged Plover . . . . .	iii	30
" <i>cannabina</i> . . . . .	ii	20	" " Sandpiper . . . . .	iii	25
" <i>Chloris</i> . . . . .	ii	11	Long-legs . . . . .	iii	30
" <i>flavirostris</i> . . . . .	ii	22	Long-neck . . . . .	iii	12
" <i>minor</i> . . . . .	ii	22	Long Pod . . . . .	i	79
" " . . . . .	ii	24	Long-shank, Black-winged . . . . .	iii	30
" <i>montana</i> . . . . .	ii	25	Long-shanks . . . . .	iii	30



	Vol.	Page		Vol.	Page
<b>Long-tailed Duck</b> . . .	iii	100	<b>Martin, Bank</b> . . .	i	141
<b>Long-tailed Mag</b> . . .	i	79	“ <b>Black</b> . . .	i	133
“ <b>“ Muffin</b> . . .	i	79	“ <b>House</b> . . .	i	139
“ <b>“ Shieldrake</b> . . .	iii	100	<b>Martin, Purple</b> . . .	i	138
“ <b>“ Pie</b> . . .	i	79	“ <b>Sand</b> . . .	i	141
“ <b>“ Tit</b> . . .	i	79	<b>Martin, Screech</b> . . .	i	133
<b>Long-tailed Titmouse</b> . . .	i	79	“ <b>Swallow</b> . . .	i	139
<b>Long Tom</b> . . .	i	79	<b>Masked Gull</b> . . .	iii	147
<b>Long-tongue</b> . . .	i	116	<b>Mate, Cuckoo's</b> . . .	i	116
<b>Loon</b> . . .	iii	108	<b>Mattageess</b> . . .	i	59
“ . . .	iii	112	<b>Mavis</b> . . .	ii	49
“ . . .	iii	114	<b>Meadow Crake</b> . . .	iii	50
“ <b>Greater</b> . . .	iii	108	“ <b>Lark</b> . . .	i	154
“ <b>Sprat</b> . . .	iii	116	“ . . .	i	158
<b>Lough Diver</b> . . .	iii	103	<b>Meadow Pipit</b> . . .	i	154
<b>Loxia Chloris</b> . . .	ii	11	<b>Meadow Titling</b> . . .	i	154
“ <i>coccothraustes</i> . . .	ii	14	<b>Mealy Redpole</b> . . .	ii	24
“ <i>curvirostra</i> . . .	ii	30	<i>Mecisura vagans</i> . . .	i	79
“ “ <i>major</i> . . .	ii	32	<b>Mediterranean Black-headed</b>		
“ <i>enucleator</i> . . .	ii	29	<b>Gull</b> . . .	iii	150
“ <i>falcirostra</i> . . .	ii	33	<i>Melizophilus provincialis</i> . . .	ii	114
“ <i>Hamburgia</i> . . .	ii	7	<b>Melodious Willow Warbler</b> . . .	ii	111
“ <i>leucoptera</i> . . .	ii	33	<b>Melodious Willow Wren</b> . . .	ii	111
“ <i>pityopsittacus</i> . . .	ii	32	<i>Melonocorypha Sibirica</i> . . .	i	163
“ <i>pyrrhula</i> . . .	ii	26	<b>Merganser, Hooded</b> . . .	iii	104
<b>Lulean Finch</b> . . .	ii	4	“ <b>Red-breasted</b> . . .	iii	105
<b>M</b>			<b>Merganser, White</b> . . .	iii	103
<i>Machetes pugnax</i> . . .	iii	33	<i>Mergoides rufina</i> . . .	iii	95
<b>Macqueen's Bustard</b> . . .	ii	152	<i>Mergulus melanoleucos</i> . . .	iii	123
<i>Macroramphus griseus</i> . . .	iii	39	<i>Mergus albellus</i> . . .	iii	103
<b>Madge</b> . . .	i	108	“ <i>castor</i> . . .	iii	106
“ <b>Owl</b> . . .	i	54	“ <i>cucullatus</i> . . .	iii	104
<b>Mag, Long-tailed</b> . . .	i	79	“ <i>fuscus</i> . . .	iii	104
<b>Magpie</b> . . .	i	108	“ <i>merganser</i> . . .	iii	106
<b>Magpie, Common</b> . . .	i	108	“ <i>minor</i> . . .	iii	116
<b>Maid, Cuckoo's</b> . . .	i	116	“ <i>serrator</i> . . .	iii	105
<b>Maize-bird, Red-winged</b> . . .	ii	38	<b>Merlin</b> . . .	i	33
<b>Mallard</b> . . .	iii	81	<i>Merops apiaster</i> . . .	i	93
<b>Manx Shearwater</b> . . .	iii	166	“ <i>chrysocephalus</i> . . .	i	93
<b>Manx Petrel</b> . . .	iii	166	“ <i>galitæus</i> . . .	i	93
“ <b>Puffin</b> . . .	iii	166	<b>Merula Iliaca</b> . . .	ii	47
<b>Mareca Americana</b> . . .	iii	88	“ <i>musica</i> . . .	ii	49
“ <i>penelope</i> . . .	iii	87	“ <i>torquata</i> . . .	ii	56
<b>Market-jew Crow</b> . . .	i	95	“ <i>viscivora</i> . . .	ii	42
<b>Marsh Blackbird</b> . . .	ii	38	“ <i>vulgaris</i> . . .	ii	53
<b>Marsh Harrier</b> . . .	i	44	<b>Messenger, Cuckoo's</b> . . .	i	116
<b>Marsh-hen</b> . . .	iii	56	<b>Mew, Sea</b> . . .	iii	156
<b>Marsh Tit</b> . . .	i	77	“ <b>Winter</b> . . .	iii	156
<b>Marsh Titmouse</b> . . .	i	77	<i>Micropus murarius</i> . . .	i	133
“ <b>Warbler</b> . . .	ii	95	<i>Milvus furcatus</i> . . .	i	24
<b>Martin</b> . . .	i	139	“ <i>ictinus</i> . . .	i	22
<b>Martin, American Purple</b> . . .	i	138	“ <i>regalis</i> . . .	i	22
			“ <i>vulgaris</i> . . .	i	22
			<b>Missel Bird</b> . . .	ii	42



	Vol.	Page		Vol.	Page
<b>Missel Thrush</b> . . . . .	ii	42	<i>Motacilla trochilus</i> . . . . .	ii	109
Misseltoe Thrush . . . . .	ii	42	“ <i>troglodytes</i> . . . . .	ii	116
Minute Dunlin . . . . .	iii	45	“ <i>Yarrellii</i> . . . . .	ii	142
Mire Drum . . . . .	iii	11	Mother Carey's Chicken . . . . .	iii	170
Moat-hen . . . . .	iii	56	<b>Mottled Owl</b> . . . . .	i	55
Mock-nightingale . . . . .	ii	97	Mountain Blackbird . . . . .	ii	56
Mokoho Bittern . . . . .	iii	13	<b>Mountain Finch</b> . . . . .	ii	4
<b>Montagu's Harrier</b> . . . . .	i	47	Mountain Linnet . . . . .	ii	25
Moor Blackbird . . . . .	ii	56	“ Pie . . . . .	i	59
“ Buzzard . . . . .	i	44	“ Sparrow . . . . .	i	177
Moor-cock . . . . .	ii	137	“ “ . . . . .	ii	7
“ -coot . . . . .	iii	56	Mouse Hawk . . . . .	i	48
“ -fowl . . . . .	ii	137	Moustache Tern . . . . .	iii	139
“ Harrier . . . . .	i	44	Muffin . . . . .	i	79
<b>Moor-hen</b> . . . . .	iii	56	“ Long-tailed . . . . .	i	79
Moor-Titling . . . . .	ii	74	Muggy . . . . .	ii	103
Mope, Blue . . . . .	i	72	Muir-fowl . . . . .	ii	137
<i>Morinellus marinus</i> . . . . .	ii	167	Mum Ruffin . . . . .	i	79
<i>Mormon fratercula</i> . . . . .	iii	124	Murdering Pie . . . . .	i	59
<i>Morpheus congener</i> . . . . .	i	11	“ “ . . . . .	i	62
Moss Creeper . . . . .	i	154	<i>Muscicapa atricapilla</i> . . . . .	i	84
<i>Motacilla acredula</i> . . . . .	ii	109	“ <i>grisola</i> . . . . .	i	87
“ <i>alba</i> . . . . .	i	142	“ <i>luctuosa</i> . . . . .	i	84
“ “ . . . . .	i	144	“ <i>muscipeta</i> . . . . .	i	84
“ <i>alpina</i> . . . . .	ii	59	<b>Mute Swan</b> . . . . .	iii	74
“ <i>arundinacea</i> . . . . .	ii	86			
“ <i>atrata</i> . . . . .	ii	72			
“ <i>atricapilla</i> . . . . .	ii	97			
“ <i>boarula</i> . . . . .	i	146			
“ <i>Brissoni</i> . . . . .	i	144			
“ <i>cinerea</i> . . . . .	i	146			
“ <i>curruca</i> . . . . .	ii	105			
“ <i>dumetorum</i> . . . . .	ii	105			
“ <i>flava</i> . . . . .	i	148			
“ “ . . . . .	i	149			
“ <i>hippolais</i> . . . . .	ii	112			
“ <i>Ischecautschia</i> . . . . .	ii	74			
“ <i>lusciniæ</i> . . . . .	ii	90			
“ <i>melanopa</i> . . . . .	i	146			
“ <i>modularis</i> . . . . .	ii	60			
“ <i>mosquitta</i> . . . . .	ii	97			
“ <i>neglecta</i> . . . . .	i	48			
“ <i>œnanthe</i> . . . . .	ii	80			
“ <i>Phœnicurus</i> . . . . .	ii	69			
“ <i>provincialis</i> . . . . .	ii	114			
“ <i>regulus</i> . . . . .	ii	119			
“ <i>rubecula</i> . . . . .	ii	62			
“ <i>rubetra</i> . . . . .	ii	76			
“ <i>rubicola</i> . . . . .	ii	74			
“ <i>Suecica</i> . . . . .	ii	67			
“ <i>sulphurea</i> . . . . .	i	146			
“ <i>sylvia</i> . . . . .	ii	103			
“ <i>sylvicola</i> . . . . .	ii	105			
“ <i>trochilus</i> . . . . .	ii	107			

## N

<i>Neuclerus furcatus</i> . . . . .	i	24
Neb, Black . . . . .	i	99
<i>Neophron percnopterus</i> . . . . .	i	3
Nettle-creeper . . . . .	ii	103
Nick-a-pecker . . . . .	i	120
Night-hawk . . . . .	i	131
<b>Night Heron</b> . . . . .	iii	10
<b>Nightingale</b> . . . . .	ii	90
“ <b>Thrush</b> . . . . .	ii	92
<b>Nightjar</b> . . . . .	i	131
“ <b>Red-necked</b> . . . . .	i	130
Nightjar, Russet-necked . . . . .	i	130
Night Raven . . . . .	iii	10
“ Warbler . . . . .	ii	86
Nine Killer . . . . .	i	62
<i>Noctua funerea</i> . . . . .	i	58
“ <i>nudipes</i> . . . . .	i	56
“ <i>nyctea</i> . . . . .	i	52
“ <i>passerina</i> . . . . .	i	56
“ <i>Tengmalmi</i> . . . . .	i	57
Nocturnal Goatsucker . . . . .	i	131
<b>Noddy</b> . . . . .	iii	144
Noddy, Black . . . . .	iii	144
“ Tern . . . . .	iii	146
Nope . . . . .	ii	24



	Vol.	Page
Norfolk Plover . . . . .	ii	155
Northern Diver . . . . .	iii	114
“ Doucker . . . . .	iii	115
“ Fulmar . . . . .	iii	165
“ Hareld . . . . .	iii	100
“ Three-toed Woodpecker . . . . .	i	125
<i>Nucifraga caryocatactes</i> . . . . .	i	140
<i>Numenius arquata</i> . . . . .	iii	19
“ <i>borealis</i> . . . . .	iii	20
“ <i>Hudsonicus</i> . . . . .	iii	20
“ <i>major</i> . . . . .	iii	19
“ <i>minor</i> . . . . .	iii	20
“ <i>pygmaeus</i> . . . . .	iii	40
“ <i>viridis</i> . . . . .	iii	17
Numidian Crane . . . . .	iii	2
Nun . . . . .	i	72
Nutcracker . . . . .	i	110
Nuthatch . . . . .	i	114
Nutjobber . . . . .	i	114
<i>Nyctihelidon Europæus</i> . . . . .	i	131
<i>Nycticorax Europæus</i> . . . . .	iii	10
<i>Nyroca ferina</i> . . . . .	iii	96
“ <i>fuligula</i> . . . . .	iii	99
“ <i>leucopthalma</i> . . . . .	iii	97
“ <i>marila</i> . . . . .	iii	98

## O

<i>Œdicnemus Bellonii</i> . . . . .	ii	155
“ <i>crepitans</i> . . . . .	ii	155
<i>Œgialitis vocifera</i> . . . . .	ii	169
<i>Œnanthe nostra tertia</i> . . . . .	ii	74
“ <i>secunda</i> . . . . .	ii	76
<i>Oidemia fusca</i> . . . . .	iii	92
“ <i>nigra</i> . . . . .	iii	93
“ <i>perspicillata</i> . . . . .	iii	94
Olivaceous Crake . . . . .	iii	53
Olive . . . . .	ii	172
<i>Onychoprion fuliginosus</i> . . . . .	iii	138
Orange-legged Hobby . . . . .	i	32
Oriole, Golden . . . . .	ii	58
<i>Oriolus galbula</i> . . . . .	ii	58
Orphean Warbler . . . . .	ii	99
Ortolan . . . . .	i	183
Ortolan Bunting . . . . .	i	183
<i>Ortygometra Baillonii</i> . . . . .	iii	54
“ <i>crex</i> . . . . .	iii	50
“ <i>porzana</i> . . . . .	iii	51
Osprey . . . . .	i	13
<i>Otis Macqueenii</i> . . . . .	ii	152
“ <i>œdicnemus</i> . . . . .	ii	155
“ <i>tarda</i> . . . . .	ii	151
“ <i>tetrax</i> . . . . .	ii	152

<i>Otus brachyotus</i> . . . . .	i	48
“ <i>vulgaris</i> . . . . .	i	49
Ouzel, Ring . . . . .	ii	56
Ouzel, Rock . . . . .	ii	56
“ Rose . . . . .	ii	34
“ “ -coloured . . . . .	ii	34
“ Water . . . . .	ii	40
Owl, Barn . . . . .	i	54
“ Brown . . . . .	i	53
“ Canada . . . . .	i	58
“ Church . . . . .	i	54
“ Churn . . . . .	i	131
Owl, Eagle . . . . .	i	50
Owl, Great . . . . .	i	50
“ “ -eared . . . . .	i	50
“ “ Horned . . . . .	i	50
“ Hawk . . . . .	i	48
Owl, Hawk . . . . .	i	58
Owl, Hissing . . . . .	i	54
Owl, Little . . . . .	i	56
Owl, Little Horned . . . . .	i	51
“ “ Night . . . . .	i	56
Owl, Long-eared . . . . .	i	49
Owl, Long-horned . . . . .	i	49
“ Madge . . . . .	i	54
Owl, Mottled . . . . .	i	55
“ Scops-eared . . . . .	i	51
Owl, Screech . . . . .	i	54
Owl, Short-eared . . . . .	i	48
Owl, Short-horned . . . . .	i	48
Owl, Snowy . . . . .	i	52
“ Tawny . . . . .	i	53
“ Tengmalm's . . . . .	i	57
Owl, Tengmalm's Night . . . . .	i	57
Owl, White . . . . .	i	54
Owl, Woodcock . . . . .	i	48
“ Yellow . . . . .	i	54
Oxeye . . . . .	i	66
“ Smaller . . . . .	i	77
Oyster-catcher . . . . .	ii	172
Oyster-catcher, Pied . . . . .	ii	172

## P

<i>Pandion haliaëtus</i> . . . . .	i	13
Parrot Crossbill . . . . .	ii	32
Parrot, German . . . . .	ii	89
Partridge . . . . .	ii	142
“ Barbary . . . . .	ii	146
Partridge, Common . . . . .	ii	142
“ French . . . . .	ii	145
“ Guernsey . . . . .	ii	145
Partridge, Red-legged . . . . .	ii	145
“ Virginian . . . . .	ii	147



	Vol.	Page		Vol.	Page
<i>Parus ater</i> . . . . .	i	69	<i>Phalacrocorax cristatus</i> . . . . .	iii	130
" <i>atricapillus</i> . . . . .	i	69	<b>Phalarope, Grey</b> . . . . .	iii	61
" " . . . . .	i	77	" <b>Red-necked</b> . . . . .	iii	62
" <i>barbatus</i> . . . . .	i	82	<i>Phalaropus fulicarius</i> . . . . .	iii	62
" <i>biarmicus</i> . . . . .	i	82	" <i>hyperboreus</i> . . . . .	iii	61
" <i>cæruleus</i> . . . . .	i	72	" " . . . . .	iii	62
" <i>caudatus</i> . . . . .	i	79	" <i>lobatus</i> . . . . .	iii	61
" <i>cristatus</i> . . . . .	i	70	" <i>platyrhynchus</i> . . . . .	iii	62
" <i>longicaudatus</i> . . . . .	i	79	<i>Phasianus Colchicus</i> . . . . .	ii	131
" <i>major</i> . . . . .	i	66	<i>Phatalea leucorodia</i> . . . . .	iii	16
" <i>palustris</i> . . . . .	i	77	" <i>nivea</i> . . . . .	iii	16
<b>Passenger Pigeon</b> . . . . .	ii	130	<b>Pheasant</b> . . . . .	ii	131
<i>Passer arundinacea minor</i> . . . . .	ii	86	<b>Pheasant, Common</b> . . . . .	ii	131
" <i>arundinaceus</i> . . . . .	i	177	" <b>Reed</b> . . . . .	i	82
" <i>domesticus</i> . . . . .	ii	9	" <b>Ring-necked</b> . . . . .	ii	131
<i>Passerine Bunting</i> . . . . .	ii	177	<i>Philomela lusciniæ</i> . . . . .	ii	90
<i>Passer montanus</i> . . . . .	ii	7	" <i>Turdoides</i> . . . . .	ii	92
<b>Pastor, Rose-coloured</b> . . . . .	ii	34	<i>Phœnicura Suecica</i> . . . . .	ii	67
<i>Pastor roseus</i> . . . . .	ii	34	" <i>ruticilla</i> . . . . .	ii	69
<b>Pectoral Sandpiper</b> . . . . .	iii	47	" <i>Tithy's</i> . . . . .	ii	72
<b>Peep</b> . . . . .	iii	45	<i>Phylloscopus superciliosus</i> . . . . .	ii	123
<b>Peep, Little</b> . . . . .	iii	45	<b>Pianet</b> . . . . .	i	108
<b>Peewit</b> . . . . .	ii	165	<i>Pica caudata</i> . . . . .	i	108
<b>Peewit Gull</b> . . . . .	iii	146	<b>Pick-a-tree</b> . . . . .	i	120
<b>Peggy Wash-dish</b> . . . . .	i	142	<i>Picoides tridactylus</i> . . . . .	i	125
<i>Pelecanus bassanus</i> . . . . .	iii	131	<i>Picus major</i> . . . . .	i	122
" <i>carlo</i> . . . . .	iii	128	" <i>martius</i> . . . . .	i	119
" <i>cristatus</i> . . . . .	iii	130	" <i>minor</i> . . . . .	i	123
" <i>graculus</i> . . . . .	iii	130	" <i>tridactylus</i> . . . . .	i	125
<i>Perdix cinerea</i> . . . . .	ii	142	" <i>varius major</i> . . . . .	i	122
" <i>coturnix</i> . . . . .	ii	149	" " <i>minor</i> . . . . .	i	123
" <i>Gibraltarica</i> . . . . .	ii	150	" " <i>tertius</i> . . . . .	i	123
" <i>petrosa</i> . . . . .	ii	146	" <i>villosus</i> . . . . .	i	124
" <i>rufa</i> . . . . .	ii	145	" <i>viridis</i> . . . . .	i	120
" <i>Virginiana</i> . . . . .	ii	147	<b>Pied Flycatcher</b> . . . . .	i	84
<b>Peregrine</b> . . . . .	i	28	<b>Pied Oyster-catcher</b> . . . . .	ii	172
<b>Peregrine-falcon</b> . . . . .	i	28	<b>Pied Wagtail</b> . . . . .	i	142
<b>Petrel, Bulwer's</b> . . . . .	iii	167	" <b>Woodpecker</b> . . . . .	i	122
<b>Petrel, Common Storm</b> . . . . .	iii	170	<b>Pie, French</b> . . . . .	i	122
<b>Petrel, Fork-tailed</b> . . . . .	iii	169	" <b>Long-tailed</b> . . . . .	i	79
<b>Petrel, Fork-tailed Storm</b> . . . . .	iii	169	" <b>Mountain</b> . . . . .	i	59
" <b>Fulmar</b> . . . . .	iii	165	" <b>Murdering</b> . . . . .	i	59
" " . . . . .	iii	165	" " . . . . .	i	62
" <b>Leach's</b> . . . . .	iii	169	" <b>Sea</b> . . . . .	ii	172
" <b>Manks</b> . . . . .	iii	166	<b>Pigeon, French</b> . . . . .	ii	165
<b>Petrel, Storm</b> . . . . .	iii	170	<b>Pigeon, Passenger</b> . . . . .	ii	130
<b>Petrel, Stormy</b> . . . . .	iii	170	" <b>Wood</b> . . . . .	ii	124
<b>Petrel, Wilson's</b> . . . . .	iii	168	<b>Pigmy Curlew</b> . . . . .	iii	40
<i>Petrocincla saxatilis</i> . . . . .	ii	52	<b>Pine Bullfinch</b> . . . . .	ii	29
<b>Pettychaps, Greater</b> . . . . .	ii	101	<b>Pine Grossbeak</b> . . . . .	ii	29
" <b>Lesser</b> . . . . .	ii	112	<b>Pink</b> . . . . .	ii	1
<b>Pe-wit</b> . . . . .	ii	165	<b>Pink-footed Goose</b> . . . . .	iii	66
<i>Phœnopterus roseus</i> . . . . .	iii	18	<b>Pinnock</b> . . . . .	i	82
<i>Phæopus arquatus</i> . . . . .	iii	20	" <b>Bearded</b> . . . . .	i	82



	Vol.	Page		Vol.	Page
Pintail . . . . .	iii	80	Pomerine Skua . . . . .	iii	162
Pintail, Common . . . . .	iii	80	Pope . . . . .	ii	26
“ Duck . . . . .	iii	80	Popinjay . . . . .	i	120
Pipit . . . . .	i	154	<i>Porphyrio rufescens</i> . . . . .	iii	50
“ Brighton . . . . .	i	153	Post Bird . . . . .	i	87
“ Lark . . . . .	i	158	Pratincole . . . . .	ii	153
Pipit, Meadow . . . . .	i	154	Pratincole, Austrian . . . . .	ii	153
“ Red-throated . . . . .	i	156	“ Collared . . . . .	ii	153
“ Richard's . . . . .	i	151	<i>Procellaria anglorum</i> . . . . .	iii	166
“ Rock . . . . .	i	160	“ <i>Bullockii</i> . . . . .	iii	169
Pipit, Shore . . . . .	i	160	“ <i>Bulwerii</i> . . . . .	iii	167
Pipit, Tawny . . . . .	i	152	“ <i>glacialis</i> . . . . .	iii	165
“ Tree . . . . .	i	158	“ <i>Leachii</i> . . . . .	iii	160
Pipit, Vinous . . . . .	i	153	“ <i>pelagica</i> . . . . .	iii	170
Pipit, Water . . . . .	i	153	“ <i>puffinus</i> . . . . .	iii	166
<i>Plectrophanes nivalis</i> . . . . .	i	171	“ <i>Wilsoni</i> . . . . .	iii	168
“ <i>lapponica</i> . . . . .	i	173	Proud-tail . . . . .	ii	16
Plover, Dotterel . . . . .	ii	159	Ptarmigan . . . . .	ii	139
Plover, Golden . . . . .	ii	157	Puckeridge . . . . .	i	131
“ Great . . . . .	ii	155	Pudding, Poke . . . . .	i	79
Plover, Green . . . . .	ii	157	Puffin . . . . .	iii	124
“ “ . . . . .	ii	165	Puffin, Manx . . . . .	iii	166
Plover, Grey . . . . .	ii	164	<i>Puffinus anglorum</i> . . . . .	iii	166
Plover, Kentish . . . . .	ii	163	Pump-borer . . . . .	i	123
Plover, Kildeer . . . . .	ii	169	Parre . . . . .	iii	48
Plover, Little Ringed . . . . .	ii	162	Purple Crested Heron . . . . .	iii	4
“ Long-legged . . . . .	iii	30	Purple Heron . . . . .	iii	4
“ Norfolk . . . . .	ii	155	“ Martin . . . . .	i	138
“ Ringed . . . . .	ii	160	“ Sandpiper . . . . .	iii	49
Plover, Sociable . . . . .	ii	170	Puttock . . . . .	i	22
Plover's-page . . . . .	iii	48	“ . . . . .	i	44
Plover, Stilt . . . . .	iii	30	<i>Pyrgita domestica</i> . . . . .	ii	9
“ Whistling . . . . .	ii	155	“ <i>montana</i> . . . . .	ii	7
“ “ . . . . .	ii	157	<i>Pyrrhocorax graculus</i> . . . . .	i	95
“ Yellow . . . . .	ii	157	<i>Pyrrhula enucator</i> . . . . .	ii	29
Pochard . . . . .	iii	96	“ <i>erythryna</i> . . . . .	ii	28
Pochard, Common . . . . .	iii	96	“ <i>vulgaris</i> . . . . .	ii	26
“ Duck . . . . .	iii	96			
“ Red-crested . . . . .	iii	95			
“ Red-headed . . . . .	iii	96			
“ Scaup . . . . .	iii	98			
“ Tufted . . . . .	iii	99			
“ Western . . . . .	iii	88			
<i>Podiceps auritus</i> . . . . .	iii	111			
“ <i>Caspicus</i> . . . . .	iii	110			
“ <i>cornutus</i> . . . . .	iii	110			
“ <i>cristatus</i> . . . . .	iii	108			
“ <i>minor</i> . . . . .	iii	112			
“ <i>obscurus</i> . . . . .	iii	110			
“ <i>rubricollis</i> . . . . .	iii	109			
Pod, Long . . . . .	i	79			
Poke Pudding . . . . .	i	79			
<i>Polysticta Stelleri</i> . . . . .	iii	88			
Pomerine Jager . . . . .	iii	162			

## Q

Quail . . . . .	ii	149
“ Andalusian . . . . .	ii	150
Quail, Common . . . . .	ii	149
“ Gibraltar . . . . .	ii	150
“ Three-toed . . . . .	ii	150
Queest . . . . .	ii	124
<i>Querquedula acuta</i> . . . . .	iii	80
“ <i>circia</i> . . . . .	iii	83
“ <i>crecca</i> . . . . .	iii	86
“ <i>Discors</i> . . . . .	iii	85

## R

Rafter . . . . .	i	87
------------------	---	----



	Vol.	Page		Vol.	Page
Rail, Dwarf . . . . .	iii	51	Redpoll . . . . .	ii	22
“ Spotted . . . . .	iii	51	Redpoll, Greater . . . . .	ii	20
Rain-bird . . . . .	i	120	“ Lesser . . . . .	ii	24
“ -crow . . . . .	i	126	“ “ . . . . .	ii	22
“ -fowl . . . . .	i	120	Redpoll, Mealy . . . . .	ii	24
“ Goose . . . . .	iii	116	Redpoll, Stony . . . . .	ii	24
<i>Rallus aquaticus</i> . . . . .	iii	55	Red-sandpiper . . . . .	iii	41
“ <i>crex</i> . . . . .	iii	50	Redshank . . . . .	iii	23
“ <i>porzana</i> . . . . .	iii	51	Redshank, Common . . . . .	iii	23
“ <i>pusillus</i> . . . . .	iii	53	Redshank, Spotted . . . . .	iii	22
Rattle-wings . . . . .	iii	102	Redstart . . . . .	ii	69
Raven . . . . .	i	97	Redstart, Black . . . . .	ii	72
Raven, Night . . . . .	iii	10	“ Blue-throated . . . . .	ii	67
“ Spotted . . . . .	iii	10	Red-tail . . . . .	ii	69
Ray's Wagtail . . . . .	i	149	“ . . . . .	ii	72
Razor-bill . . . . .	iii	126	Red-throated Diver . . . . .	iii	116
<i>Recurvirostra avocetta</i> . . . . .	iii	29	“ Pipit . . . . .	i	156
“ <i>avocetta</i> . . . . .	iii	29	Redwing . . . . .	ii	47
Red-backed Shrike . . . . .	i	62	Red-winged Blackbird . . . . .	ii	38
Red-billed Heron . . . . .	iii	8	“ Maize-bird . . . . .	ii	38
Redbreast . . . . .	ii	62	Red-winged Starling . . . . .	ii	38
Red-breasted Bernicle . . . . .	iii	69	Reed Fauvette . . . . .	ii	84
Red-breasted Flycatcher . . . . .	i	86	“ Pheasant . . . . .	i	82
Red-breasted Goosander . . . . .	iii	105	Reed Warbler . . . . .	ii	86
Red-breasted Goose . . . . .	iii	69	Reed Wren . . . . .	ii	86
Red-breasted Linnet . . . . .	ii	20	Reeve . . . . .	iii	33
Red-breasted Merganser . . . . .	iii	105	<i>Regulus atricapillus</i> . . . . .	ii	119
Red-breasted Shoveler . . . . .	iii	78	“ <i>cristatus</i> . . . . .	ii	119
“ Snipe . . . . .	iii	32	Regulus, Dalmatian . . . . .	ii	123
“ “ . . . . .	iii	39	<i>Regulus ignicapillus</i> . . . . .	ii	122
Redbreast, Robin . . . . .	ii	62	“ <i>modestus</i> . . . . .	ii	123
Red-cap . . . . .	ii	16	“ <i>non-cristatus major</i> . . . . .	ii	107
Red-crested Duck . . . . .	iii	95	“ <i>trochilus</i> . . . . .	ii	109
“ Pochard . . . . .	iii	95	“ <i>vulgaris</i> . . . . .	ii	119
Red-crested Whistling Duck . . . . .	iii	95	<i>Rhyncaspis clypeata</i> . . . . .	iii	78
Red Duck . . . . .	iii	97	Richardson's Arctic Gull . . . . .	iii	163
Red-fronted Swallow . . . . .	i	136	Richardson's Skua . . . . .	iii	163
Red Grouse . . . . .	ii	137	Richard's Pipit . . . . .	i	151
Red Godwit . . . . .	iii	32	Rinding-bird . . . . .	i	116
Red-headed Pochard . . . . .	iii	96	Ring Dove . . . . .	ii	124
Red-leg . . . . .	iii	23	Ringed Dotterel . . . . .	ii	160
Red-legged Crow . . . . .	i	95	“ Guillemot . . . . .	iii	120
“ Gull . . . . .	iii	148	Ringed Plover . . . . .	ii	160
“ Heron . . . . .	iii	9	Ring-necked Pheasant . . . . .	ii	131
“ Horseman . . . . .	iii	23	Ring Ouzel . . . . .	ii	56
“ Jackdaw . . . . .	i	95	Ring Thrush . . . . .	ii	56
Red-legged Partridge . . . . .	ii	145	<i>Rissa Brunnichii</i> . . . . .	iii	152
Red-legged Sandpiper . . . . .	iii	23	Robin . . . . .	ii	62
Red-necked Coot-foot . . . . .	iii	62	“ Blue-throated . . . . .	ii	67
Red-necked Grebe . . . . .	iii	109	Robinet . . . . .	ii	62
Red-necked Lobe-foot . . . . .	iii	62	Robin Redbreast . . . . .	ii	62
Red-necked Nightjar . . . . .	i	130	Rock Dove . . . . .	ii	128
“ Phalarope . . . . .	iii	62	Rockier . . . . .	ii	128
			Rock Lark . . . . .	i	160



	Vol.	Page
Rock Ouzel . . . . .	ii	56
Rock Pipit . . . . .	ii	160
“ Thrush . . . . .	ii	52
Roller . . . . .	i	89
Roller, Garrulous . . . . .	i	89
Rook . . . . .	i	103
Roseate Tern . . . . .	iii	135
Rose-coloured Ouzel . . . . .	ii	34
Rose-coloured Pastor . . . . .	ii	34
Rose-coloured Starling . . . . .	ii	34
“ Linnet . . . . .	ii	20
“ Ouzel . . . . .	ii	34
Rotche . . . . .	iii	123
Rotche, Common . . . . .	iii	123
Rough-legged Buzzard . . . . .	i	18
Royston Crow . . . . .	i	101
<i>Rubetra anglicana</i> . . . . .	i	84
Ruddock . . . . .	ii	62
Ruddy Shelduck . . . . .	iii	77
Ruff . . . . .	iii	33
Ruff, Fighting . . . . .	iii	33
Ruffin, Mum . . . . .	i	79
Rufous-backed Egret . . . . .	iii	8
“ Bittern . . . . .	iii	12
Rufous Warbler . . . . .	ii	95
Runner . . . . .	iii	55
Russet-necked Nightjar . . . . .	i	130
Rustic Bunting . . . . .	i	179
<i>Ruticilla phœnicurus</i> . . . . .	ii	69

## S

Sabine's Gull . . . . .	iii	145
<i>Salicaria arundinacea</i> . . . . .	ii	86
“ <i>galactotes</i> . . . . .	ii	95
“ <i>locustella</i> . . . . .	ii	82
“ <i>luscinioides</i> . . . . .	ii	83
“ <i>phragmitis</i> . . . . .	ii	84
“ <i>Turdoides</i> . . . . .	ii	93
Sandcock . . . . .	iii	23
Sanderling . . . . .	ii	171
Sand Grouse . . . . .	ii	141
Sand Lark . . . . .	ii	160
“ “ . . . . .	iii	26
“ Lavrock . . . . .	ii	160
“ “ . . . . .	iii	26
Sand Martin . . . . .	i	141
Sandpiper, Ash-coloured . . . . .	iii	41
Sandpiper, Bartram's . . . . .	iii	42
Sandpiper, Black . . . . .	iii	49
Sandpiper, Broad-billed . . . . .	iii	44
Sandpiper, Brown . . . . .	iii	45
Sandpiper, Buff-breasted . . . . .	iii	43
“ Common . . . . .	iii	26

	Vol.	Page
Sandpiper, Curlew . . . . .	iii	40
Sandpiper, Dunlin . . . . .	iii	48
“ Dusky . . . . .	iii	22
“ Equestrian . . . . .	iii	33
“ Gambet . . . . .	iii	23
Sandpiper, Green . . . . .	iii	24
Sandpiper, Greenwich . . . . .	iii	33
“ Grey . . . . .	ii	164
“ Hebridial . . . . .	ii	167
“ Lapwing . . . . .	ii	165
“ Little . . . . .	iii	45
“ “ . . . . .	iii	46
“ Long-legged . . . . .	iii	25
Sandpiper, Pectoral . . . . .	iii	47
“ Purple . . . . .	iii	49
Sandpiper, Red-legged . . . . .	iii	23
“ Redshank . . . . .	iii	23
Sandpiper, Schinz's . . . . .	iii	46
Sandpiper, Sea . . . . .	iii	49
“ Selninger . . . . .	iii	49
“ Shore . . . . .	iii	33
“ Spotted . . . . .	iii	26
Sandpiper, Spotted . . . . .	iii	27
Sandpiper, Striated . . . . .	iii	23
“ Temminck's . . . . .	iii	46
“ Whistling . . . . .	iii	24
“ Wood . . . . .	iii	24
Sandpiper, Wood . . . . .	iii	25
Sandpiper, Yellow-legged . . . . .	iii	33
“ Yellow-shanked . . . . .	iii	43
Sandwich Tern . . . . .	iii	133
Savi's Warbler . . . . .	ii	83
<i>Saxicola ananthe</i> . . . . .	ii	80
“ <i>deserti</i> . . . . .	ii	79
“ <i>isabellina</i> . . . . .	ii	78
“ <i>rubetra</i> . . . . .	ii	76
“ <i>rubicola</i> . . . . .	ii	74
“ <i>stapazina</i> . . . . .	ii	78
Scare-crow . . . . .	i	101
Scarlet Bullfinch . . . . .	ii	28
Scarlet Grossbeak . . . . .	ii	28
Scaup . . . . .	iii	98
Scaup Duck . . . . .	iii	98
“ Pochard . . . . .	iii	98
Schinz's Sandpiper . . . . .	iii	46
Sclavonian Grebe . . . . .	iii	110
Scobby . . . . .	ii	1
<i>Scolopax agrocephala</i> . . . . .	iii	31
“ <i>arquata</i> . . . . .	iii	19
“ <i>Belgica</i> . . . . .	iii	31
“ <i>borealis</i> . . . . .	iii	20
“ <i>calidris</i> . . . . .	iii	23
“ <i>canescens</i> . . . . .	iii	28



	Vol.	Page		Vol.	Page
<i>Scolopax flavipes</i>	iii	43	Shearwater Petrel	iii	166
“ <i>fuscus</i>	iii	22	<b>Shelduck</b>	iii	76
“ <i>gallinago</i>	iii	36	“ <b>Ruddy</b>	iii	77
“ <i>gallinaria</i>	iii	36	Shell-apple	ii	1
“ <i>gallinula</i>	iii	38	“	ii	30
“ <i>glottis</i>	iii	28	Shieldrake	iii	76
“ <i>grisea</i>	iii	39	“ Common	iii	76
“ <i>Hudsonica</i>	iii	31	“ Long-tailed	iii	100
“ <i>Lapponica</i>	iii	31	Shilfa	ii	1
“ <i>leucophæa</i>	iii	32	<b>Shore Lark</b>	i	160
“ <i>limosa</i>	iii	31	Shore Pipit	i	160
“ <i>major</i>	iii	35	“ Sandpiper	iii	33
“ <i>noveboracensis</i>	iii	39	<b>Short-eared Owl</b>	i	48
“ “	iii	32	Short-heeled Field Lark	i	158
“ <i>phaeopus</i>	iii	20	“ -horned Owl	i	48
“ <i>pygmæus</i>	iii	40	<b>Short-toed Lark</b>	i	164
“ <i>rusticola</i>	iii	34	<b>Shoveler</b>	iii	78
“ <i>totanus</i>	iii	22	Shoveler, Blue-winged	iii	78
“ “	iii	23	“ Common	iii	78
Scooper	iii	29	“ Duck	iii	78
Scooping Avocet	iii	29	“ Red-breasted	iii	78
<i>Scops Aldrovandi</i>	i	51	Shreek	i	59
<b>Scops-eared Owl</b>	i	51	Shrieker	iii	31
<b>Scoter</b>	iii	93	Shrike	i	59
Scoter, Black	iii	93	“ Ash-coloured	i	59
“ Common	iii	93	“ Cinereous	i	59
<b>Scoter, Turf</b>	iii	94	“ -cock	ii	42
“ <b>Velvet</b>	iii	94	“ Great	i	59
<i>Scotornis trimaculatus</i>	i	130	“ “ Grey	i	59
Scout	iii	117	<b>Shrike, Grey</b>	i	59
Scraper, Common	iii	121	“ Lesser Grey	i	61
Screamer	i	133	“ Red-backed	i	62
Screech	i	133	Shrike, Wood	i	64
“ Martin	i	133	“ Woodchat	i	64
“ Owl	i	54	Shrite	ii	42
Sea Hen	iii	117	Shuffle-wing	ii	60
Sea Lark	i	160	Siberian Goose	iii	69
“ “	iii	48	Silk-tail	i	113
“ Mew	iii	156	Silver-winged Black Tern	iii	143
Sea-parrot	iii	124	Silvery Gull	iii	158
Sea Pie	ii	172	<b>Siskin</b>	ii	18
“ Sandpiper	iii	49	<i>Sitta Europæa</i>	i	114
“ Snipe	iii	48	Sit-ye-down	i	66
“ Titling	i	160	Skeeling Goose	iii	76
Sedge Bird	ii	84	Skelgoose	iii	76
<b>Sedge Warbler</b>	ii	84	Skelly	ii	1
Sedge Wren	ii	84	Skite	i	180
Selninger Sandpiper	iii	49	<b>Skua</b>	iii	161
Senegal Bittern	iii	9	Skua, Arctic	iii	164
<b>Serin Finch</b>	ii	13	“ Buffon's	iii	164
Shag	iii	130	“ Common	iii	161
“ Crested	iii	130	“ Gull	iii	161
Sharp-tailed Duck	iii	100	<b>Skua, Long-tailed</b>	iii	164
<b>Shearwater, Manx</b>	iii	166	“ Pomerine	iii	162



	Vol.	Page		Vol.	Page
Skua, Richardson's . . .	iii	163	Speckled Diver . . .	iii	116
Sky Lark . . .	i	167	"    Loon . . .	iii	115
Small Doucker . . .	iii	112	Spink . . .	ii	1
Smaller Oxye . . .	i	77	<i>Spinus linaria</i> . . .	ii	22
Smew . . .	iii	103	Spoonbill . . .	iii	16
Snake-bird . . .	i	116	Spoonbill, White . . .	iii	16
Snipe . . .	iii	36	Spotted Crake . . .	iii	51
Snipe, Black-headed . . .	iii	22	"    Eagle . . .	i	11
Snipe, Brown . . .	iii	39	"    Flycatcher . . .	i	87
Snipe, Common . . .	iii	36	Spotted Gallinule . . .	iii	51
"    Courland . . .	iii	22	"    Guillemot . . .	iii	121
"    Double . . .	iii	35	"    Rail . . .	iii	51
"    Dusky . . .	iii	22	"    Raven . . .	iii	10
Snipe, Great . . .	iii	35	Spotted Redshank . . .	iii	22
Snipe, Grey . . .	iii	39	"    Sandpiper . . .	iii	26
"    Half . . .	iii	38	Spotted Sandpiper . . .	iii	27
Snipe, Jack . . .	iii	38	"    Snipe . . .	iii	22
Snipe, Jadreka . . .	iii	31	"    Water-hen . . .	iii	51
"    Least . . .	iii	45	Sprat Loon . . .	iii	116
"    " . . .	iii	48	Squacco Heron . . .	iii	9
"    Pool . . .	iii	23	Squaiotta Heron . . .	iii	9
"    Red-breasted . . .	iii	32	<i>Squatarola cinerea</i> . . .	ii	164
"    " . . .	iii	39	Squealer . . .	i	133
"    Sea . . .	iii	48	Stannel Hawk . . .	i	35
"    Solitary . . .	iii	35	Stare . . .	ii	35
"    Spotted . . .	iii	22	"    Collared . . .	ii	59
"    Summer . . .	iii	26	"    Common . . .	ii	35
"    Whole . . .	iii	36	Starling . . .	ii	35
Snite . . .	iii	36	Starling, Common . . .	ii	35
Snow Bunting . . .	i	171	Starling, Red-winged . . .	ii	38
Snow Flake . . .	i	171	Starling, Rose-coloured . . .	ii	34
"    Fleck . . .	i	171	St. Cuthbert's Duck . . .	iii	89
"    Fowl . . .	i	171	Steller's Eider . . .	iii	88
Snow Goose . . .	iii	69	<i>Stercorarius cephus</i> . . .	iii	164
Snowy Owl . . .	i	52	" <i>longicaudus</i> . . .	iii	164
Sociable Plover . . .	ii	170	<i>Sterna Anglica</i> . . .	iii	140
Solan Gannet . . .	iii	131	" <i>arctica</i> . . .	iii	137
"    Goose . . .	iii	131	" <i>Boysii</i> . . .	iii	133
Solitary Snipe . . .	iii	35	" <i>canliaca</i> . . .	iii	133
"    Thrush . . .	ii	35	" <i>Caspia</i> . . .	iii	132
<i>Somateria dispar</i> . . .	iii	88	" <i>Dougalli</i> . . .	iii	135
" <i>mollissima</i> . . .	iii	89	" <i>fissipes</i> . . .	iii	142
" <i>spectabilis</i> . . .	iii	91	" <i>fuliginosa</i> . . .	iii	138
Song Thrush . . .	ii	49	" <i>hirundo</i> . . .	iii	136
Sooty Tern . . .	iii	138	" <i>leucopareia</i> . . .	iii	139
Sparrow . . .	ii	9	" <i>leucoptera</i> . . .	iii	143
Sparrow, Common . . .	ii	9	" <i>minuta</i> . . .	iii	141
Sparrow-hawk . . .	i	41	" <i>nævia</i> . . .	iii	142
Sparrow, House . . .	ii	9	" <i>nigra</i> . . .	iii	142
"    Mountain . . .	i	177	" <i>stolida</i> . . .	iii	144
"    " . . .	ii	7	Stilt . . .	iii	30
Sparrow, Tree . . .	ii	7	Stilt, Black-winged . . .	iii	30
Sparrow, Water . . .	i	177	"    Plover . . .	iii	30
<i>Spathulea clypeata</i> . . .	iii	78	Stint . . .	iii	45



	Vol.	Page		Vol.	Page
Stint . . . . .	iii	48	Swallow, Common . . . . .	i	136
“ Little . . . . .	iii	45	“ Martin . . . . .	i	139
Stint, Temminck's . . . . .	iii	46	“ Red-fronted . . . . .	i	136
Stock Dove . . . . .	ii	126	“ Swift . . . . .	i	133
Stone-chacker . . . . .	ii	80	Swallow-tailed Kite . . . . .	i	24
Stonechat . . . . .	ii	74	Swallow, Window . . . . .	i	139
Stonechatter . . . . .	ii	74	Swamp Blackbird . . . . .	ii	38
Stoneclink . . . . .	ii	74	Swan, Bewick's . . . . .	iii	73
Stone Curlew . . . . .	ii	155	Swan, Hooping . . . . .	iii	72
Stonegall . . . . .	i	35	Swan, Mute . . . . .	iii	74
Stone-smith . . . . .	ii	74	Swan, Tame . . . . .	iii	74
Stony Redpole . . . . .	ii	24	“ Whistling . . . . .	iii	72
Stork, Black . . . . .	iii	15	Swift . . . . .	i	133
“ White . . . . .	iii	14	“ Alpine . . . . .	i	135
Storm-cock . . . . .	ii	42	Swift, Common . . . . .	i	133
Storm Petrel . . . . .	iii	170	“ Swallow . . . . .	i	133
Storm Petrel, Common . . . . .	iii	170	“ White-bellied . . . . .	i	135
Stormy Petrel . . . . .	iii	170	Swinepipe . . . . .	ii	47
<i>Streptopelia interpres</i> . . . . .	ii	167	<i>Sylvia ænanthe</i> . . . . .	ii	80
Striated Sandpiper . . . . .	iii	23	“ <i>arundinacea</i> . . . . .	ii	86
Striped Diver . . . . .	iii	116	“ <i>atricapilla</i> . . . . .	ii	97
<i>Strix aluco</i> . . . . .	i	53	“ <i>cinerea</i> . . . . .	ii	103
“ <i>asio</i> . . . . .	i	55	“ <i>curruca</i> . . . . .	ii	105
“ <i>brachyotus</i> . . . . .	i	48	“ <i>Dartfordiensis</i> . . . . .	ii	114
“ <i>bubo</i> . . . . .	i	50	“ <i>dumetorum</i> . . . . .	ii	105
“ <i>dasyptus</i> . . . . .	i	56	“ <i>Gibraltariensis</i> . . . . .	ii	72
“ <i>flammea</i> . . . . .	i	54	“ <i>grisea</i> . . . . .	ii	99
“ <i>funerea</i> . . . . .	i	57	“ <i>hippolais</i> . . . . .	ii	111
“ “ . . . . .	i	58	“ “ . . . . .	ii	112
“ <i>giu</i> . . . . .	i	51	“ <i>hortensis</i> . . . . .	ii	101
“ <i>nudipes</i> . . . . .	i	56	“ <i>ignicapillus</i> . . . . .	ii	122
“ <i>nyctea</i> . . . . .	i	52	“ <i>locustella</i> . . . . .	ii	82
“ <i>otus</i> . . . . .	i	49	“ <i>lusciniæ</i> . . . . .	ii	90
“ <i>passerina</i> . . . . .	i	56	“ <i>luscinioides</i> . . . . .	ii	83
“ <i>scops</i> . . . . .	i	51	“ <i>modularis</i> . . . . .	ii	60
“ <i>stridula</i> . . . . .	i	53	“ <i>nisoria</i> . . . . .	ii	96
“ <i>Tengmalmi</i> . . . . .	i	57	“ <i>Orphea</i> . . . . .	ii	99
“ <i>ulula</i> . . . . .	i	48	“ <i>phænicurus</i> . . . . .	ii	69
“ <i>zorca</i> . . . . .	i	51	“ <i>phragmitis</i> . . . . .	ii	84
<i>Sturnia funerea</i> . . . . .	i	58	“ <i>provincialis</i> . . . . .	ii	114
<i>Sturnis vulgaris</i> . . . . .	ii	35	“ <i>regulus</i> . . . . .	ii	119
<i>Sturnus cinclus</i> . . . . .	ii	40	“ <i>rubecula</i> . . . . .	ii	62
“ <i>collaris</i> . . . . .	ii	59	“ <i>rubetra</i> . . . . .	ii	76
“ <i>moritanicus</i> . . . . .	ii	59	“ <i>rubicola</i> . . . . .	ii	74
“ <i>prædatorius</i> . . . . .	ii	58	“ <i>rufa</i> . . . . .	ii	112
Sucker, Cherry . . . . .	i	87	“ <i>salicaria</i> . . . . .	ii	84
<i>Sula alba</i> . . . . .	iii	131	“ <i>sibilatrix</i> . . . . .	ii	107
“ <i>bassana</i> . . . . .	iii	131	“ <i>Suecica</i> . . . . .	ii	67
Summer Snipe . . . . .	iii	26	“ <i>sylvicola</i> . . . . .	ii	107
Surf Scoter . . . . .	iii	94	“ <i>sylvicola</i> . . . . .	ii	105
<i>Surnia nyctea</i> . . . . .	i	52	“ <i>Tithys</i> . . . . .	ii	72
Swabian Bittern . . . . .	iii	9	“ <i>trochilus</i> . . . . .	ii	109
Swallow . . . . .	i	136	“ <i>troglodytes</i> . . . . .	ii	116
Swallow, Chimney . . . . .	i	136	“ <i>Turdoides</i> . . . . .	ii	92



<i>Syrnium Aluco</i> . . .	Vol. Page
<i>Syrrhaptes paradoxus</i> . . .	i 53
	ii 141

## T

<i>Tadorna Bellonii</i> . . .	iii 76	<i>Thalassidroma Wilsoni</i> . . .	iii 168
“ <i>rutila</i> . . .	iii 77	Thick-billed Guillemot . . .	iii 119
“ <i>vulpanser</i> . . .	iii 76	“ -knee . . .	ii 155
Tame Swan . . .	iii 74	“ “ Common . . .	ii 155
<i>Tantalus falcinellus</i> . . .	iii 17	“ -kneed Bustard . . .	ii 155
Tarrock . . .	iii 117	Thistle-finch . . .	ii 16
Tattler . . .	iii 43	Three-toed Quail . . .	ii 150
“ Bartram's . . .	iii 42	Three-toed Woodpecker . . .	i 125
Tawny Bunting . . .	i 171	Thrush . . .	ii 49
“ Lark . . .	i 152	Thrush, Common . . .	ii 49
Tawny Owl . . .	i 53	“ Holme . . .	ii 42
“ Pipit . . .	i 152	Thrush, Missel . . .	ii 42
Teal . . .	iii 86	Thrush, Misseltoe . . .	ii 42
“ Blue-winged . . .	iii 85	Thrush Nightingale . . .	ii 92
Teal, Common . . .	iii 86	“ Rock . . .	ii 52
“ Garganey . . .	iii 83	Thrush, Solitary . . .	ii 35
Teazer . . .	iii 164	“ Song . . .	ii 49
Temminck's Dunlin . . .	iii 46	Thrush, White's . . .	ii 51
“ Sandpiper . . .	iii 46	Thrush, Wind . . .	ii 47
Temminck's Stint . . .	iii 46	Throstle . . .	ii 49
Tengmalm's Night Owl . . .	i 57	Tippet Grebe . . .	iii 108
Tengmalm's Owl . . .	i 57	Tit, Bearded . . .	i 82
Tern, Arctic . . .	iii 137	“ Blue . . .	i 72
“ Black . . .	iii 142	“ Bottle . . .	i 79
“ Caspian . . .	iii 132	“ Cole . . .	i 69
“ Common . . .	iii 136	“ Crested . . .	i 70
“ Gull-billed . . .	iii 140	“ Great . . .	i 66
“ Lesser . . .	iii 141	Tithy's Redstart . . .	ii 72
Tern, Little . . .	iii 141	Titlark . . .	i 154
“ Moustache . . .	iii 139	Titling . . .	i 154
“ Noddy . . .	iii 144	“ Meadow . . .	i 154
Tern, Roseate . . .	iii 135	“ Sea . . .	i 160
“ Sandwich . . .	iii 133	“ Tree . . .	i 158
Tern, Silver-winged Black . . .	iii 143	Tit, Long-tailed . . .	i 79
Tern, Sooty . . .	iii 138	“ Marsh . . .	i 77
“ Whiskered . . .	iii 139	Titmouse, Bearded . . .	i 82
“ White-winged Black . . .	iii 143	“ Blue . . .	i 72
<i>Tetrao colurnix</i> . . .	ii 149	“ Cole . . .	i 69
“ <i>lagopus</i> . . .	ii 139	“ Crested . . .	i 70
“ <i>perdix</i> . . .	ii 142	“ Great . . .	i 66
“ <i>rufus</i> . . .	ii 145	“ Long-tailed . . .	i 79
“ <i>Scoticus</i> . . .	ii 137	“ Marsh . . .	i 77
“ <i>tetrix</i> . . .	ii 135	Tom, Bottle . . .	i 79
“ <i>urogallus</i> . . .	ii 133	“ Collier . . .	i 66
Te-wit . . .	ii 165	“ Long . . .	i 79
<i>Thalassidroma Bullockii</i> . . .	iii 169	Tomtit . . .	i 72
“ <i>Bulwerii</i> . . .	iii 167	“ Great Black-headed . . .	i 66
“ <i>Leachii</i> . . .	iii 168	Tongue-bird . . .	i 116
“ <i>pelagica</i> . . .	iii 170	<i>Tolanus Bartramia</i> . . .	iii 42
		“ <i>Bartramii</i> . . .	iii 42
		“ <i>calidris</i> . . .	iii 23
		“ <i>cinereus</i> . . .	iii 33
		“ <i>flavipes</i> . . .	iii 43
		“ <i>fuscus</i> . . .	iii 22







	Vol.	Page		Vol.	Page
Wagtail, Ray's . . . .	i	149	Wheatear, Black-throated	ii	78
“ Water . . . .	i	142	“ Desert . . . .	ii	79
“ White . . . .	i	142	“ Isabelline . . . .	ii	78
Wagtail, White . . . .	i	144	Wheel-bird . . . .	i	131
Wagtail, Winter . . . .	i	142	Whimbrel . . . .	iii	20
“ “ . . . .	i	146	Whimbrel, Curlew . . . .	iii	20
“ Yellow . . . .	i	146	Whinchat . . . .	ii	76
“ “ . . . .	i	148	Whin Linnet . . . .	ii	20
Wagtail, Yellow . . . .	i	149	Whiskered Tern . . . .	iii	139
Wall Creeper . . . .	i	118	Whiskey John . . . .	i	62
Warbler, Alpine . . . .	ii	59	Whistling Plover . . . .	ii	155
“ Aquatic . . . .	ii	89	“ “ . . . .	ii	157
“ Barred . . . .	ii	96	“ Sandpiper . . . .	iii	24
Warbler, Blackcap . . . .	ii	97	“ Swan . . . .	iii	72
“ Blue-throated . . . .	ii	67	White-bellied Swift . . . .	i	135
Warbler, Dartford . . . .	ii	114	“ -eyed Duck . . . .	iii	97
Warbler, Epicurean . . . .	i	84	White-fronted Goose . . . .	iii	67
Warbler, Garden . . . .	ii	101	White-fronted Wild Goose . . . .	iii	67
“ Grasshopper . . . .	ii	82	White-headed Goosander . . . .	iii	103
“ Great Sedge . . . .	ii	93	“ “ Harpy . . . .	i	44
“ Marsh . . . .	ii	95	“ Merganser . . . .	iii	103
“ Melodious Willow . . . .	ii	111	“ Nun . . . .	iii	103
Warbler, Night . . . .	ii	86	White Owl . . . .	i	54
Warbler, Orphean . . . .	ii	99	White Spoonbill . . . .	iii	16
“ Reed . . . .	ii	86	White's Thrush . . . .	ii	47
“ Rufous . . . .	ii	95	White Stork . . . .	iii	14
“ Savi's . . . .	ii	83	White-tail . . . .	ii	80
“ Sedge . . . .	ii	84	Whitethroat . . . .	ii	103
“ Willow . . . .	ii	109	Whitethroat, Common . . . .	ii	103
“ Wood . . . .	ii	107	Whitethroat, Lesser . . . .	ii	105
Warbler, Yellow . . . .	ii	107	White Wagtail . . . .	i	142
“ “ . . . .	ii	109	White Wagtail . . . .	i	144
Warbler, Yellow-browed . . . .	ii	122	“ -winged Black Tern . . . .	iii	143
Wash-dish, Peggy . . . .	i	142	“ “ Crossbill . . . .	ii	33
Washer, Dish . . . .	i	142	“ “ Lark . . . .	i	163
Water-crake . . . .	iii	51	White-winged Silvery Gull . . . .	iii	155
“ Crow . . . .	ii	40	Whitterick . . . .	iii	19
“ -hen . . . .	iii	56	Whittle . . . .	i	120
“ -hen . . . .	iii	55	Whitwall . . . .	i	122
“ “ Spotted . . . .	iii	51	Whole Snipe . . . .	iii	36
“ Ouzel . . . .	ii	40	Wigeon . . . .	iii	87
Water Pipit . . . .	i	154	“ American . . . .	iii	88
Water-rail . . . .	iii	51	Wigeon, Common . . . .	iii	87
Water Sparrow . . . .	i	177	Wild Duck . . . .	iii	81
“ Wagtail . . . .	i	142	Wild Goose . . . .	iii	63
Waxen Chatterer . . . .	i	113	“ “ . . . .	iii	65
Waxwing . . . .	i	113	Willow . . . .	iii	117
“ Bohemian . . . .	i	113	Willow-biter . . . .	i	77
Western Duck . . . .	iii	88	Willow Warbler . . . .	ii	109
“ Eider . . . .	iii	88	Willow Wren . . . .	ii	109
“ Pochard . . . .	iii	88	Wilson's Petrel . . . .	iii	168
Whaap . . . .	iii	19	Windhover . . . .	i	35
“ . . . .	iii	20	Window Swallow . . . .	i	139
Wheatear . . . .	ii	80	Wind Thrush . . . .	ii	47



	Vol.	Page		Vol.	Page
Winter Fauvette . . .	ii	60	Wren, Reed . . .	ii	86
" Mew . . .	iii	156	" Sedge . . .	ii	84
" Wagtail . . .	i	142	" Willow . . .	ii	109
" " . . .	i	146	" Wood . . .	ii	107
Witwall . . .	i	122	" Yellow . . .	ii	107
Woodchat . . .	i	64	Wryneck . . .	i	116
Woodchat Shrike . . .	i	64			
Woodcock . . .	iii	34	<b>X</b>		
Woodcock Owl . . .	i	48	<i>Xema atricilla</i> . . .	iii	151
Woodcracker . . .	i	114	" <i>minutus</i> . . .	iii	146
Wood Grouse . . .	ii	133	" <i>Sabini</i> . . .	iii	145
Wood Lark . . .	i	165			
Woodnacker . . .	i	122	<b>Y</b>		
Woodpecker Barred . . .	i	123	Yaffer . . .	i	120
Woodpecker, Black . . .	i	119	Yaffle . . .	i	120
Woodpecker, French . . .	i	122	Yappingal . . .	i	120
" Great Black . . .	i	119	Yarwhelp . . .	iii	31
" " " and White . . .	i	122	Yarwhip . . .	iii	31
Woodpecker Great Spotted . . .	i	122	Yeldring . . .	i	180
Woodpecker, Greater Spotted . . .	i	122	Yeldrock . . .	i	180
Woodpecker Green . . .	i	120	Yellow-billed Cuckoo . . .	i	126
" Hairy . . .	i	124	" -browed Warbler . . .	ii	123
Woodpecker, Large Green . . .	i	129	Yellow Bunting . . .	i	180
" Least . . .	i	123	Yellow-hammer . . .	i	180
Woodpecker, Lesser Spotted . . .	i	123	Yellow-hammer, Black-throated . . .	i	182
Woodpecker, Little Black and White . . .	i	123	" " French . . .	i	182
" " French . . .	i	123	" -legged Gull . . .	iii	157
" Pied . . .	i	122	" " Sandpiper . . .	iii	33
Woodpecker, Three-toed . . .	i	125	" Owl . . .	i	54
Woodpecker, Three-toed Northern . . .	i	125	" Plover . . .	ii	157
Woodpie . . .	i	122	" -shanked Sandpiper . . .	iii	43
Wood Pigeon . . .	ii	124	Yellow-shanks . . .	iii	43
Wood Sandpiper . . .	iii	24	Yellow-throated Bee-eater . . .	i	93
Wood Sandpiper . . .	iii	25	" Wagtail . . .	i	146
Wood Shrike . . .	i	64	" " . . .	i	148
Woodspite . . .	i	120	Yellow Wagtail . . .	i	149
Woodwall . . .	i	122	Yellow Warbler . . .	ii	107
Wood Warbler . . .	ii	107	" " . . .	ii	109
Wood Wren . . .	ii	107	" Willow Wren . . .	ii	107
Wren . . .	ii	116	" Yeldring . . .	i	180
Wren, Common . . .	ii	116	" Yite . . .	i	180
" Fire-crested . . .	ii	122	" Yoldring . . .	i	180
" Furze . . .	ii	114	" Yowley . . .	i	180
" Golden-crested . . .	ii	119	Yelper . . .	iii	29
" " -crowned . . .	ii	119	Yite, Yellow . . .	i	180
" Green . . .	ii	107	Yoit . . .	i	180
" Kitty . . .	ii	116	Yoldring, Yellow . . .	i	180
" Larger Willow . . .	ii	107	Yolkwing, " . . .	i	180
" Least Willow . . .	ii	112	" Yowley, " . . .	i	180
" Melodious Willow . . .	ii	111	<i>Funx torquilla</i> . . .	i	111




## Z

	Vol. Page		Vol. Page
<i>Zapornia Baillonii</i> .	iii 54	<i>Zapornia pusilla</i> .	iii 53

END OF VOLUME III.















RETURN TO the circulation desk of any  
University of California Library  
or to the

NORTHERN REGIONAL LIBRARY FACILITY  
Bldg. 400, Richmond Field Station  
University of California  
Richmond, CA 94804-4698

---

ALL BOOKS MAY BE RECALLED AFTER 7 DAYS  
2-month loans may be renewed by calling  
(415) 642-6233

1-year loans may be recharged by bringing books  
to NRLF

Renewals and recharges may be made 4 days  
prior to due date

---

DUE AS STAMPED BELOW

---

**JUL 08 1989**

---

---

---

---

---

---

---

---

---

---



Morris, F.O

A natural history of  
the nests and eggs of  
British birds.

1892

V.3

BIOLOGY  
LIBRARY  
G

APR 1 1938

4-6

held - APR 9 1938

983265 QL675

M6

1892

V.3  
BIOLOGY  
LIBRARY  
G

THE UNIVERSITY OF CALIFORNIA LIBRARY



